


Hosted by:


Freshwater Mollusk Conservation Society

Mollusks . People . Streams



FMCS

Mollusk Health & Disease Workshop
March 13-15, 2018



11th Biennial Workshop

2018 Workshop Organizers

Diane Waller
U.S. Geological Survey
Upper Midwest Environmental Sciences Center

Megan Bradley & Nathan Eckert
U.S. Fish and Wildlife Service
Genoa National Fish Hatchery

Symposium Planning Committee

Lisie Kitchel, Jeremy Tiemann & Steve Ahlstedt- Sponsorships
Shelly Bartsch- Poster section
Heidi Dunn- Mollusk Die-off Section Coordination
Corey Puzach- Laboratory Section Coordination and Local Arrangements
Emily Grossman- Management of Exchequer
Louise LaVictoire- Program layout and coordination
Greg Cope-Presenter selection

2017 FMCS Officers

<i>President</i>	<i>Past President</i>
Heidi Dunn	Teresa Newton
Ecological Specialists, Inc. O'Fallon, MO	US Geological Survey La Crosse, WI
<i>President Elect</i>	<i>Secretary</i>
Jeremy Tiemann	Janet Clayton
Illinois Natural History Survey Champaign, IL	West Virginia Division of Natural Resources Elkins, WV
<i>Treasurer</i>	
Emily Robbins	
Ecological Specialists, Inc. O'Fallon, MO	

FMCS Standing Committees and Their Chairs/Co-chairs

If you are interested in participating in committee activities, please contact one of the appropriate chairs.

Awards

W. Gregory Cope - North Carolina State University greg_cope@ncsu.edu
Teresa Newton - Upper Midwest Environ. Science Center
newton@usgs.gov
Emy Monroe - Midwest Fisheries Center emy_monroe@fws.gov

Environmental Quality & Affairs

Steve McMurray - Missouri Dept. of Conservation
stephen.mcmurray@mdc.mo.gov

Braven Beaty - The Nature Conservancy bbeaty@tnc.org

Gastropod Status and Distribution

Nathan Whelan - Auburn University nwhelan@auburn.edu

Genetics

David J. Berg - Miami University bergdj@miamioh.edu
Dave Zanatta - Central Michigan University zanat1d@cmich.edu

Guidelines and Techniques

Ryan Schwegman - EnviroScience, Inc.
RSchwegman@EnviroScienceInc.com

Lisie Kitchel- Wisconsin Dept. Nat. Resources lisie.kitchel@wisconsin.gov

Information Exchange Newsletter

John Jenkinson - Clinton, Tennessee jjjenkinson@hotmail.com

Information Exchange Journal

W. Gregory Cope - North Carolina State University greg_cope@ncsu.edu
Wendell R. Haag - U.S. Forest Service whaag@fs.fed.us

David J. Berg - Miami University bergdj@miamioh.edu

Mussel Status and Distribution

Arthur E. Bogan – North Carolina State Museum of Natural Sciences
arthur.bogan@ncdenr.gov

John L. Harris - Arkansas State University omibob1@gmail.com

Nominations

Leroy Koch - U.S. Fish and Wildlife Service leroy_koch@fws.gov

Outreach


Jennifer Archambault – N.C. State University jmarcham@ncsu.edu

Propagation, Restoration, & Introduction

Rachael Hoch – N.C. Wildlife Resources Comm.
rachael.hoch@ncwildlife.org

Tim Lane- Virginia Dept.of Game & Inland Fisheries
tim.lane@dgif.virginia.gov

Symposium Jeremy Tiemann - Illinois Natural History Survey
jtiemann@illinois.edu



Radisson – Ground Level

The Freshwater Mollusk Conservation Society (FMCS) is dedicated to the conservation and advocacy of freshwater mollusks for public education and the conservation science of freshwater mollusks, North America's most imperiled fauna.

River Level Sponsors (≥\$1000)



DEPARTMENT OF
**GAME & INLAND
FISHERIES**
CONSERVE. CONNECT. PROTECT.

Stream Level Sponsors (>\$500 – \$999)



Eddy Level Sponsors (\$100 – \$499)



Schedule

Monday 3/12	Tuesday 3/13	Wednesday 3/14	Thursday 3/15
	Breakfast 7:30-8:00 am	Breakfast 7:30-8:00 am	Breakfast 7:30-8:00 am
	Welcome	<p>Small Group Sessions 8:00-11:30</p> <p>Lab session: group 1 (pink heelsplitters- red dot on nametag) 8:00am bus leaves</p> <p>Risk characterization session: group 2 (green floaters-green dot on nametag)- Radisson ballroom</p> <p>Die-offs and kills session: group 3 (yellow sandshell-yellow dot on nametag)- Radisson ballroom</p>	<p>Small Group Sessions 8:00-11:30</p> <p>Lab session: group 3 (yellow sandshell- yellow dot on nametag) 8:00am bus leaves</p> <p>Risk characterization session: group 1 (pink heelsplitters- red dot on nametag)- Radisson ballroom</p> <p>Die-offs and kills session: group 2 (green floaters-green dot on nametag)- Radisson ballroom</p>
	8:00-8:10		
	The Uncertain State of Molluscan Health		
	Dr. Greg Cope and Diane Waller 8:10-8:40		
	Back to the Drawing Board: Assessing Causes of Freshwater Mussel Declines		
	Wendell Haag 8:45-9:30		
	Viruses of freshwater mussels (and other species): expanding the invertebrate “virosphere” in the pursuit of mussel health and conservation		
	Tony Goldberg 9:30-10:00		
	Break 10:00-10:15		
	An Overview of the Bacteriological Examination of Freshwater Mussels		
	Eric Leis 10:15-10:45		
	Are Parasites and Diseases Contributing to the Decline of Freshwater Mussels (Bivalvia, Unionida)?		
	Andrew McElwain 10:45-11:15		
	Perspective from the Marine Realm: Contemporary Challenges and Approaches to Bivalve Mollusc Health Management		
	Ryan Carnegie 11:15-12:00		
	Plated Lunch 12:00-1:30	Groups 1 & 2 lunch provided, Group 3 lunch on own	Group 3 lunch provided, Groups 1 & 2 lunch on own

Monday 3/12	Tuesday 3/13	Wednesday 3/14	Thursday 3/15
	<p>Overview of Health Assessment Tools in Native Freshwater Mollusks</p> <p>Dr. Greg Cope and Dr. Teresa Newton 1:30-1:45</p> <p>Omics</p> <p>Ieva Roznere 1:50-2:05</p> <p>Use of Condition Indexes for Mussel Health Assessment</p> <p>Serena Ciparis 2:10-2:25</p> <p>Hemolymph chemistry profiles and fatty acid analysis as tools for evaluating freshwater mussel health</p> <p>Andrea Fritts 2:30-2:45</p> <p>Influence of water and sediment on the digestive gland microbiome in the Alabama rainbow (<i>Villosa nebulosa</i>)</p> <p>Cova Arias 2:50-3:10</p> <p>Break 3:10-3:25</p> <p>Disease Risk Analysis – Applications for the Management of Fresh Water Mussels</p> <p>Tiffany Wolf 3:25-3:45</p> <p>Do the Infectious Diseases of Dreissenids Represent a Threat to North American Unionid Populations?</p> <p>Daniel Molloy 3:50-4:05</p>	<p>Small Group 1:00-4:30</p> <p>Lab session: group 2 (green floaters-green dot on nametag) 12:00pm bus leaves</p> <p>Risk characterization session: group 3 (yellow sandshell-yellow dot on nametag)- Radisson ballroom</p> <p>Die-offs and kills session: group 1 (pink heelsplitters- red dot on nametag)- Radisson ballroom</p>	<p>Tour of Genoa National Fish Hatchery, Bus Leaves 1pm 1:00-5:00</p>
<p>Registration Open 4:00-9:00</p>	<p>Invasive snails and their potential to serve as hosts for parasites in the Midwest</p> <p>Greg Sandland 4:10-4:25</p> <p>Factors Related to Growth Inhibition in Juvenile Mussels Exposed to Ambient Stream Conditions</p> <p>Wendell Haag 4:30-4:45</p> <p>1999 Ohio River Mollusk Kill Assessment: The Gastropod Study</p> <p>Janet Clayton 4:50-5:05</p> <p>Freshwater mussel die-offs: insights from a compilation of known cases.</p> <p>Jordan Richards 5:10-5:30</p> <p>Poster Session and Social 7:00-9:00</p>	<p>FMCS Board Meeting 5:00-7:00</p> <p>Social and Jam Session 7:00-9:00</p>	

Speaker Biographies:

Ryan Carnegie is an Associate Research Scientist at Virginia Institute of Marine Science, Department of Environmental & Aquatic Animal Health. He received a BA in Biology from Rutgers University, MA in Marine Science from VIMS, and PhD in Marine Biology from the University of Maine. His research interests include parasitology and invertebrate pathology; shellfish genetics and molluscan population biology and host-parasite interactions.

Dr. W. Gregory Cope is a William Neal Reynolds Distinguished Professor in the Department of Applied Ecology at NC State University. He also serves as Department Extension Leader and Coordinator of the NC Agromedicine Institute. Dr. Cope's research interests are in aquatic toxicology, ecology, and physiology, and in the transport, fate, and effects of aquatic pollutants and other human-mediated stressors on freshwater mollusks and fish. His research utilizes sentinel aquatic organisms, biomarkers of exposure, effect, or susceptibility, or alternative toxicological models from which linkages to environmental and human health are evaluated. Major areas of research focus include the assessment of pesticides, persistent organochlorine contaminants, and metals in surface waters, the effects of waterborne and sediment-associated contaminants on fish and native mollusks in inland waterways, the efficacy of constructed wetlands and other Best Management Practices for reducing non-point source pollution from urban (e.g., polycyclic aromatic hydrocarbons) and agricultural (e.g., nutrients) watersheds, and the effects of contaminant availability and cycling on imperiled species. Dr. Cope is a Charter Member of FMCS and has held numerous officer and committee positions in the Society, including President and Chair of the Awards Committee, and currently serves as Co-Editor of the Society's Journal, *Freshwater Mollusk Biology and Conservation*.

Andrea Fritts is a biologist with the USGS Upper Midwest Environmental Sciences Center. Her past research has included aspects of early life history, sclerochronology, and physiology of freshwater mussels, while her current research uses stable isotopes and fatty acids to study food web interactions. Andrea received her PhD from the University of Georgia.

Tony Goldberg is a professor of epidemiology at the University of Wisconsin-Madison, Department of Pathobiological Sciences School of Veterinary Medicine. He received a BA from Amherst College, a PhD from Harvard University, and DVM from University of Illinois. His research focuses on the ecology, epidemiology and evolution of infectious disease, combining field and laboratory studies to understand how pathogens in dynamic ecosystems are transmitted among hosts, across complex landscapes, and over time.

Wendell Haag is a Research Fisheries Biologist with the U.S. Forest Service, Southern Research Station. His research on freshwater mussels spans 30 years and has explored a variety of topics, including life history, fish-host relationships, age and growth, population and assemblage dynamics, biogeography, and conservation issues. He has authored over 50 peer-reviewed papers, book chapters, and books, including *North American Freshwater Mussels: Natural History, Ecology, and Conservation* (2012). His current research focuses on identifying causes and mechanisms of mussel declines. He has served as co-editor of *Freshwater Mollusk Biology and Conservation* since 2012 and was Associate Editor of *Freshwater Science* from 2009 to 2015.

Susan Knowles is a diagnostic pathologist at the USGS National Wildlife Health Center, Madison, WI. She received a BA from Bucknell University, DVM from Virginia-Maryland Regional College of Veterinary Medicine, and PhD from the University of Georgia. Her research interests include wildlife and aquatic animal pathology, including bivalves and novel virus discovery.

Eric Leis is a fish biologist at the UFWS La Crosse Fish Health Center. He studies parasite infections in fish and the morphological and genetic identification of these parasites. He also develops cell lines from fish and amphibians applied in the fields of virology and toxicology.

Ieva Roznere is a postdoctoral researcher at The Ohio State University. She is a physiological ecologist researching how environmental stress affects freshwater mussel metabolism using metabolomic and transcriptomic techniques.

Heidi Dunn is a malacologist, with EcoAnalyst (formerly Ecological Specialists). She holds a BS in Wildlife Science from Purdue University and a MS in Biology from Southern Illinois University at Edwardsville, IL. She was introduced to freshwater mussels in 1978 while working as a summer technician with USFWS, and started sampling and working with other malacologists in the Mississippi River basin in the 1980s. She has been working with these animals ever since and never ceases to be amazed at their diversity, beauty, and behavior. She has conducted over 600 mussel study designs, surveys, relocations, Biological Assessments, and workshops since founding Ecological Specialists in 1990. She has sampled throughout the interior basin, on the Atlantic coast and down to Texas. She is also currently an instructor for USFWS Conservation of Freshwater Mussel Biology at the National Conservation Training Center in Shepherdstown, WV and President of the Freshwater Mollusk Conservation Society.

Dr. Teresa J. Newton is a fishery biologist with the U.S. Geological Survey's Upper Midwest Environmental Sciences Center in La Crosse, WI. She received a B.S. in Biology in 1985 from Central Michigan University, a M.S. in Biology in 1987 from Tennessee Technological University, and a Ph.D. in Fisheries Biology and Toxicology in 1990 from Iowa State University. She has been employed at the USGS lab since 1990 where her research interests focus on the conservation and ecology of freshwater mussels, a group of benthic animals in which 70% of the North American species are considered threatened. Mussels are keystone species in many rivers and their catastrophic decline may lead to the decline of other faunal groups and the alteration of ecosystem processes. She uses a combination of comparative and experimental approaches to understand factors affecting the distribution and abundance of freshwater mussels, identify biochemical and physiological indicators of stress, and determine the roles that mussels play in large river food webs. Collectively, these studies provide information to help resource managers design effective restoration and conservation strategies for native freshwater mussels.

Andrew McElwain is a visiting assistant professor at The State University of New York, Oswego. He received a BS from Westfield State College, MS from Middle Tennessee State University, and PhD from Auburn University. He uses light and scanning electron microscopy to study tissue structure of fishes and freshwater mussels to better understand how cells, tissues and organs functions. He is especially interested in histopathology of parasitic infections.

Dan Molloy is a freshwater invertebrate pathologist conducting fundamental research on the diseases of North American and Eurasian populations of *Dreissena* spp. With a passion for environmental protection, his research also includes developing target-specific, pathogen-based methods for managing these mussels. He is, for example, the patent inventor of using the bacterium *Pseudomonas fluorescens* strain CL0145A as a highly-specific control agent against these mussels – the approach used in the commercial product Zequanox[®]. A Research Scientist and Adjunct Professor at the State University of New York Great Lakes Center at Buffalo, he also maintains affiliations with the State University of New York at Albany and the University of Illinois Natural History Survey at Urbana-Champaign. In addition, he directs Molloy & Associates, LLC – a firm specializing in developing credible prevention, detection, rapid response, and eradication/control programs for *Dreissena* spp.

Dr. Nick Phelps is the Director of the Minnesota Aquatic Invasive Species Research Center and Assistant Professor in the Fisheries, Wildlife and Conservation Biology Department at the University of Minnesota. His research focuses on emerging threats to the health and sustainability of aquatic ecosystems at the intersection of humans, animals and the environment. He aims to identify threats, mitigate risks and develop long-term science-based management solutions to meet the needs of all stakeholders involved.

Tiffany Wolf is an assistant professor in the Veterinary Population Medicine Department of the University of Minnesota (UMN) College of Veterinary Medicine; she also holds an appointment as an Associate Fellow at the UMN Institute on the Environment. With a background in wildlife epidemiology and ecosystem health, she maintains a broad interest in understanding diseases of wildlife populations at the interface of humans, animals, and the environment. She is particularly interested in the dynamics of infectious diseases in multi-host pathogen systems and optimizing epidemiological, ecological, and molecular approaches to assess changes in health and mitigate disease impacts on populations. Her work entails working across scientific disciplines to find solutions to complex health problems. Current projects include understanding the role of health in a declining moose populations, *Parelaphostrongylus tenuis* and *Mycobacterium tuberculosis* complex transmission in multihost systems, human-primate disease transmission, wildlife disease surveillance, and contaminants of emerging concern in aquatic ecosystems.

TUESDAY MARCH 13, 2018

7:00 – 8:00 a.m.	CONTINENTAL BREAKFAST
8:00 – 8:10 a.m.	WELCOME AND ANNOUNCEMENTS
8:10 – 8:40 a.m.	<p>THE UNCERTAIN STATE OF MOLLUSCAN HEALTH Diane L. Waller & Dr. W. Gregory Cope</p>
	<p>Assessing the health of native freshwater mussels and snails, whether at the population or organismal level, is difficult because of the lack of established benchmarks by which to judge what is considered suitable or normal. Moreover, the lack of development of techniques, assays, and diagnostic tools that are critical in assessing the relative health and condition of mollusks has been hampered by concerted research efforts and limited financial resources. In this presentation, we outline the goals and objectives of the 2018 FMCS Freshwater Mollusk Health Workshop and summarize the history of progress in this topic area since the first workshop on freshwater mussel health and die-offs was held in 1986. Additionally, we will provide an overview of some of the existing and emerging health assessment tools that will be discussed in more detail throughout the workshop. We assert that a world-wide effort should be made in the areas of freshwater molluscan health and disease diagnosis and treatment, similar to those that currently exist for fish health and disease. As a greater number of state and federal fish hatcheries are undertaking mollusk propagation and culture programs for restoration and other efforts, it will be critical to understand the health of mussels held in captivity as well as those in the wild.</p>
8:45 – 9:30 a.m.	<p>BACK TO THE DRAWING BOARD: ASSESSING CAUSES OF FRESHWATER MUSSEL DECLINES Wendell Haag</p>
	<p>Freshwater mussel declines are on par with declines in pollinators, amphibians, bats, and other organisms but have received much less attention. Many mussel declines share the following characteristics: 1) commenced about 1970–1990; 2) affect entire assemblages; 3) curtailment of recruitment, followed by gradual faunal loss; 4) occurrence in otherwise intact streams; 5) upstream progression in some cases. Conventional explanations have three main characteristics: 1) focus on long-term, cumulative factors, such as sedimentation; 2) include multiple, unrelated factors, invoked to varying degrees; 3) causal factors often vague (e.g., “poor land use practices”) and mostly untested. These explanations do not correspond well to characteristics of recent declines. Rather, the similarity in characteristics and timing of declines suggests a single, severe factor specific to mussels acting rapidly across a large area. Vague and untested assumptions about causes provide little guidance for conservation and may have directed resources unproductively. I propose that mussel conservation be refocused along the following three points. 1) Largely discard conventional wisdom about causes of declines until it is evaluated critically. 2) Build a body of case studies that provide rigorous evaluation of the success of conservation strategies. Evaluation should emphasize short-term response variables such as individual condition or performance and deemphasize assemblage- and population-level responses, which occur slowly. 3) Increased emphasis on research and adaptive management that can identify specific causes of mussel declines. Previously ignored, but potentially widespread factors such as disease and the effects of invasive <i>Corbicula fluminea</i> should be revisited.</p>

9:30 – 10:00 a.m.	VIRUSES OF FRESHWATER MUSSELS (AND OTHER SPECIES): EXPANDING THE INVERTEBRATE “VIROSPHERE” IN THE PURSUIT OF MUSSEL HEALTH AND CONSERVATION Tony Goldberg, Chris Dunn, Jordan Richard, Diane Waller, Eric Leis & Joel Putnam
--------------------------	--

Recent advances in metagenomic technology have revealed a remarkable diversity of heretofore unknown viruses in invertebrates and their environments. As this “virosphere” continues to expand, it will become increasingly important to understand what the presence or a virus (or multiple viruses) means for population health and conservation. This talk presents results of several “virus hunting” efforts that have used these emerging technologies to identify the causes of population declines across multiple species. Data show that many species have a “normal” or “typical” virome consisting of a tractable number of endemic and apparently non-pathogenic viruses. As a consequence, defining the “baseline” virome of a species is critical for documenting changes, such as introduction of a new virus or enhanced replication of an endemic virus. Challenges to these approaches include the management of “big data,” choosing appropriate bioinformatic methods, distinguishing “real” viruses from viruses that have integrated into host genomes, and interpreting the clinical significance of results. In the case of freshwater mussels, these problems are amplified by the extreme divergence of the novel viruses we have found from their known relatives, as well as limited comparative data from other species. Nevertheless, this overall approach has great potential to elucidate the role of viruses in freshwater mussel health and conservation. A recent die-off event in the Clinch River, Tennessee, will be used to illustrate these principles.

Resources:

[Redefining the Invertebrate Virosphere available at https://molluskconservation.org/Library/pdf/MARCH2018/nature20167.pdf with password FMCS workshop 2018](https://molluskconservation.org/Library/pdf/MARCH2018/nature20167.pdf)

Morning Break 10:00 – 10:15 a.m.

10:15 - 10:45 a.m.	BACTERIOLOGY Eric Leis
---------------------------	----------------------------------

The diagnosis of bacterial disease in freshwater mussels is greatly hindered by a lack of information regarding the microbial communities associated with these animals. This makes identification of the causative agent(s) during mussel mortality events difficult, if not nearly impossible. A necessary initial step is the documentation of the normal bacterial flora. This basic information is essential to provide much needed insight into these assemblages; potentially resulting in the identification of pathogens. While there have been a few studies focused on the collection of this data, seasonally examining Unionid populations throughout geographically widespread locations will be necessary to properly evaluate bacterial species associated with mussels. The identification of these communities will also require careful examination as there are many aspects of sampling freshwater mussels which can make the interpretation of results challenging. Bacteria are a typical dietary component for many mussels and their presence may simply represent a food source, not necessarily indicating either a symbiotic or pathogenic relationship. Additionally, the close contact of the internal organs to the aquatic environment, which increases the likelihood for sample contamination, can further complicate interpretation. In an effort to examine the bacterial communities associated with Unionids in the Upper Mississippi River (UMR) basin, we cultured bacteria from the hemolymph of several mussel populations throughout the 2017 field season. Molecular methods targeting the 16S rRNA gene were used to identify the isolates. Hemolymph was selected due to its relative disconnect with the aquatic environment and role in the circulatory system. The most commonly cultured bacteria from mussels in the UMR were identified as belonging to the genus *Bacillus*. In October 2017, additional bacterial isolates were identified from hemolymph samples acquired during a mortality event of pheasantshells (*Actinonaias pectorosa*) on the Clinch River near Kyles Ford, Tennessee. These isolates were then compared to the bacteria obtained from mussels in the UMR which revealed that both mussel populations shared many common bacteria as well as some differences. Future bacteriology investigations will improve our understanding of the microbial communities associated with freshwater mussels, allowing us to identify those bacterial species capable of causing disease, and ultimately lead to improved Unionid restoration efforts.

10:45 - 11:15 a.m.

ARE PARASITES AND DISEASES CONTRIBUTING TO THE DECLINE OF FRESHWATER MUSSELS (BIVALVIA, UNIONIDA)?
Andrew McElwain

Freshwater mussels or pearly mussels (Mollusca: Bivalvia: Unionida) comprise approximately 843 nominal species in six families worldwide, but nearly half of these species are imperiled. Mussels have a unique life history starting with a parasitic larva that attaches to skin or gill of a fish for several days before transforming into a sedentary suspension feeder in the benthos. Gravid females brood their larvae in their gills for several weeks or months before releasing them, and many species have intriguing host attraction strategies to ensure the attachment of their larvae. This life history renders mussels vulnerable to toxic contaminants and the loss of benthic sediments, which are significant causes of mussel declines. Etiological agents have the potential to be contributing factors, but parasites and diseases of freshwater mussels remain understudied relative to marine bivalves. Mussels may host a wide variety of eukaryotic organisms including ciliates (Ciliophora), trematodes (Platyhelminthes: Aspidogastrea, and Digenea), roundworms (Nematoda), “moss animals” or bryozoans (Ectoprocta, Entoprocta), oligochaetes (Annelida: Oligochaeta), leeches (Annelida: Hirudinida), mites (Arthropoda: Acari), copepods (Arthropoda: Copepoda), insects (Arthropoda: Insecta), and fish eggs (Chordata: Actinopterygii). Some of these groups contain species that are considered parasitic because they appear to injure their host through attachment, feeding or otherwise invade tissue (e.g., digeneans). The literature also contains many observations of small organisms living in the mantle cavity or attached to the shell in which the relationship between the two parties is unclear (e.g., oligochaetes). Regarding diseases, there are a few published accounts of mass mortality events known as “die-offs” and “mussel kills”, and only one of these documents gross and histopathological changes to tissues. Additionally, there are some observations of tumors or neoplasms, and several accounts of shell anomalies such as parasite-induced pearl formation, and deformities such as infoldings and protuberances of the periostracum, erosion to the nacre and or periostracum, and misshapened shells. The cause of diseases and shell deformities has rarely been established and more often subject to speculation. With the above in mind, we are facing a significant biodiversity crisis and it is the intention of this review to highlight the need to understand the health of wild and captive mussels.

11:15 - 12:00 a.m.

PERSPECTIVE FROM THE MARINE REALM: CONTEMPORARY CHALLENGES AND APPROACHES TO BIVALVE MOLLUSC HEALTH MANAGEMENT
Ryan B. Carnegie

Understanding and managing health and disease in freshwater mussels can be informed by long experience in shellfish pathology in marine systems. Marine bivalve molluscs are infected by myriad microbial pathogens, the most notable of which have usually been protozoan but with bacteria also contributing and viral pathogens causing acute contemporary concern. Several of these pathogens, representing major protozoan lineages, are notifiable to the World Organisation for Animal Health. For many of these pathogens, devastating initial effects on shellfish populations and economies dependent on them were followed by success in control, for example through natural selection or selective breeding for disease resistance. Enlightened management includes strong elements of conservation, such as the use of sanctuaries from harvest to promote natural selection for resistance evolution in oyster populations. Yet many challenges remain. Disease surveillance in many areas for many mollusc species is inadequate, detection of emerging bacterial and viral pathogens in particular is problematic, basic life history knowledge for many pathogens is lacking. Difficulty in meeting these challenges is particularly troublesome given the rapid growth of shellfish aquaculture, which may promote disease emergence by creating new interactions between pathogens and host populations that have no adaptation to them.

As aquaculture production expands around the world, managing the health of aquacultured populations, including preventing undesired disease interactions with wild animal populations, will increasingly be a challenge. Despite best efforts to maintain biosecurity, widening commerce alters distributions of pathogens and hosts, producing outbreaks of established diseases and the emergence of new ones as noted above. Anthropogenic impacts on the marine environment including ocean warming contribute to shifting host and pathogen distributions and introduce other possible influences on disease processes, from harmful algal blooms and ocean acidification to organic pollutants and microplastic contamination—all of which can affect freshwater environments as well. In this evolving environmental landscape, the path to more effective management of animal health in aquaculture is not fully clear. Application of contemporary genetic methods for pathogen detection and discovery such as next-generation sequencing has garnered recent attention as part of the solution. Regardless of available diagnostics, however, disease occurrence remains a product of host and environmental factors in addition to those associated with the pathogen. Thus meeting the challenges of protecting aquaculture systems in a changing world will require incorporation of broader expertise into aquatic animal health, including chemistry and toxicology, physiology, genetics, oceanography, and ecology in addition to traditional areas of parasitology, microbiology, and virology to better understand more complex and multi-factorial etiologies. Capacity to explore disease-associated factors in controlled experiments will remain essential. Fundamental to maintaining capacity for understanding disease in aquatic ecosystems will be societal commitment to broad-based science programs in support of the goal of aquaculture to support a growing human population in the 21st century, but also to maintain the ecological integrity of aquatic ecosystems worldwide.

Plated Lunch 12:00 – 1:30 p.m.

1:30 – 1:45 p.m.	OVERVIEW OF HEALTH ASSESSMENT TOOLS IN NATIVE FRESHWATER MOLLUSKS Dr. W. Gregory Cope & Dr. Teresa J. Newton
<p>The study of freshwater mollusks has advanced substantially in many topical areas (e.g., propagation, toxicology, ecology) over the past 25 years. Although these advances have propelled the science of mollusk conservation forward, areas such as physiology, immunology, and basic biochemistry have received much less attention, largely due to limited financial resources and few investigators conducting research in these areas. The deficit of advances in these particular areas has hampered the development of tests, assays, and diagnostic tools that are critical in assessing the relative health and condition of mollusks. To address the continued questions about mollusk health, as well as to assess the new issues of disease, unexplained die-offs, and emerging contaminant exposures, improved tools and techniques are needed. Here we provide an overview of the major categories of existing tools (i.e., biomarkers) used to assess mollusk health and make inference to those requiring additional research and development. For our purposes, a biomarker is a change in a biological response at the molecular, cellular, biochemical, physiological, or behavioral level that can be related to exposure, effects, or susceptibility. Although highly useful and informative in assessing relative condition of mollusks and other organisms, connections between biomarker responses and tangible outcomes for characterizing “good health” will be imperative to future mollusk conservation and protection efforts.</p> <p>Resources: Newton and Cope, Chapter 10 Biomarker Responses of Unionid Mussels to Environmental Contaminants pdf available at https://molluskconservation.org/Library/pdf/MARCH2018/Newton_Cope-Ch10.pdf with password FMCS_workshop_2018</p>	
1:50 – 2:05 p.m.	OMICS Ieva Roznere
<p>What is omics? How has it been used in research on mollusks? Can these techniques be applied to improve conservation and management efforts? This presentation will address these questions with a focus on transcriptomics and metabolomics. Omics refers to fields of study that aim to detect and characterize a group of biological molecules in an organism, such as genes (as in the field of genomics), transcripts (transcriptomics), proteins (proteomics), and metabolites (metabolomics). Transcriptomics is the study of transcripts, the subset of genes that are being expressed at a certain time period. Metabolomics is the study of metabolites, the intermediates and products of metabolism that are produced by enzymatic chemical reactions. Because gene expression and metabolite production are closely associated with environmental conditions, studying changes in these biological molecules is especially helpful in understanding how organisms react to different environmental stressors. For example, transcriptomics and metabolomics have been used to identify stress-responses of freshwater mussels to heat, drought, pollutants, and translocation. The different fields of omics can be used as health assessment tools to provide snapshots of molluscan physiology at various levels of biological organization. Although still in its infancy, the application of omics techniques in the management of species of concern holds great potential for advancing the field of conservation biology.</p>	

<p>2:10 – 2:25 p.m.</p>	<p>USE OF CONDITION INDEX FOR MUSSEL HEALTH ASSESSMENT Serena Ciparis, Ty Stephenson, Garrett Rhyne & Susan Lingenfelter</p>
<p>Energy storage and allocation are critical metrics of freshwater mussel health due to direct relationships to survival, growth, and reproductive capability. Condition indices (CIs) assess energy storage, or ‘fatness’, by normalizing weight to size. Crosby and Gale (1990) recommended a standard method for determining bivalve CI, where standard CI = dry soft tissue weight (g) *1000/ internal shell cavity capacity (g). However, this method is rarely applied in studies of freshwater mussels. We compared eight potential methods for determining CI of adult <i>Lampsilis fasciola</i> exposed to clean (pond) water, clean (pond) sediment, elevated major ion concentrations (simulated Powell River water) and coal-contaminated sediment (Powell River sediment) in a 2x2 factorial design. Calculated CIs included two methods for live mussels (ratios of weight to length and weight to volume) and six methods for dissected mussels (ratios of tissue wet weight and tissue dry weight to shell length, cavity volume, and cavity weight). Glycogen content of mantle tissue and activities of enzymes associated with the glutathione antioxidant system in digestive gland tissue were also determined. There were no statistically significant differences between treatments for CIs calculated using live mussels or using tissue wet weight of dissected mussels. However, all CIs calculated using tissue dry weight indicated statistically significant differences between water treatments, with the standard CI yielding the most precise measurements. Glycogen content of mantle tissue was also significantly different between water treatments for male mussels only. We hypothesize that exposure to major ions caused mussels to utilize energy stores to compensate for osmotic stress, with males utilizing glycogen and females utilizing lipids from oocyte resorption. Differences between treatments would not have been detected without dissection, emphasizing the utility of the standard CI and the need for correlation with non-lethal endpoints. Results of other studies will be presented for comparison.</p> <p>Resources: A Review and Evaluation of Bivalve Condition Index Methodologies with a Suggested Standard Method is available at https://molluskconservation.org/Library/pdf/MARCH2018/Crosby_Gale_1990.pdf with password FMCS_workshop_2018</p>	
<p>2:30 – 2:45 p.m.</p>	<p>HEMOLYMPH CHEMISTRY PROFILES AND FATTY ACID ANALYSIS AS TOOLS FOR EVALUATING FRESHWATER MUSSEL HEALTH Andrea Fritts, Robert Bringolf, Brent Knights, Lynn Bartsch & Michelle Bartsch</p>
<p>The future of effective programs for monitoring the health of imperiled mussel species is reliant on the development of non-lethal biomonitoring techniques. Potential biomarkers include hemolymph chemistry profiles and fatty acid analysis. Studies indicate that extracting hemolymph from adductor muscles or extracting tissue biopsies from foot or mantle tissue does not adversely affect the long term survival of large-bodied mussel species (e.g. <i>Elliptio crassidens</i>, <i>Quadrula quadrula</i>), but may negatively affect the survival of small-bodied species (e.g. <i>Villosa vibex</i>). A growing number of researchers are using hemolymph profiles as a means of assessing mussel response to stressors such as elevated temperature, dewatering, and elevated carbon dioxide exposure (pCO_2). Hemolymph ion concentrations, glucose, and transaminase enzymes have all been documented to exhibit responses to various stress exposures. Hemolymph biomarker responses may vary under different types of stressors as well as among individuals of different size, sex, and species; therefore additional work is needed to better characterize the optimal ranges for hemolymph parameters. Fatty acid analysis is an emerging tool within the world of mollusk health. Recent research has evaluated the relationship between survival, growth, and fatty acid profiles of mussels over a longitudinal riverine gradient with varying phytoplankton assemblages. Fatty acids are also being used to evaluate freshwater mussel status over a large spatial gradient in the Midwest with various stressors, including the presence of bigheaded carps. Future work should strive to identify optimal concentrations of different fatty acids for freshwater mussels and to evaluate how fatty acids respond to controlled stress exposures.</p>	

<p>2:50 – 3:10 p.m.</p>	<p>INFLUENCE OF WATER AND SEDIMENT ON THE DIGESTIVE GLAND MICROBIOME IN THE ALABAMA RAINBOW (<i>VILLOSA NEBULOSA</i>) Alison K. Aceves, Cova R. Arias, Paul Johnson, and Stephen A. Bullard</p>
<p>Freshwater mussels (Bivalvia: Unionidae) are the most imperiled faunal group in North America. Alabama harbors the highest biodiversity of freshwater mussels in the world and currently leads restoration efforts across the Southeast. In collaboration with the Alabama Department of Conservation and Natural Resources, we are investigating the role of bacterial communities, using 16S rRNA gene sequencing, found in cultured and in-stream mussels and how factors such as water and sediment influence these communities. We collected a total of 22 Alabama rainbows (<i>Villosa nebulosa</i>) from the Alabama Aquatic Biodiversity Center reared from two different wild stocks (Shoal Creek = 5 and Flannigan Creek = 5) and collected in-stream (Shoal Creek = 10, Flannigan Creek = 2). Overall, the microbial communities between hatchery-reared and wild mussels were significantly different. Rearing environment exerted a stronger effect than population although differences were noted between mussels collected from Shoal Creek and Flannigan Creek. The most dominant phylum from cultured <i>V. nebulosa</i> was Tenericutes (>43.3%), followed by Proteobacteria (25.3%), Firmicutes (12.6%), Cyanobacteria (8.3%), and Bacteroidetes (3.1%). The top five phyla from in-stream Shoal Creek <i>V. nebulosa</i> were Tenericutes (36.4%), Chlamydiae (18%), Proteobacteria (17.3%), Bacteroidetes (9.2%), and Firmicutes (7.1%); however the top five phyla from in-stream Flannigan Creek <i>V. nebulosa</i> were Proteobacteria (44.5%), Bacteroidetes (17.3%), Firmicutes (10.2%), Acidobacteria (6.5%), and Actinobacteria (6.1%). Analysis of similarity percentages showed that the most abundant genus in cultured Shoal Creek mussels was <i>Ureaplasma</i> sp. (41.4%), whereas, in cultured Flannigan Creek mussels <i>Spiroplasma</i> sp. (23.5%) was the most abundant genus, both belonging to the phylum Tenericutes. Results from multidimensional scaling and analysis of similarity showed that water and sediment have no influence on the diversity of gut bacterial communities found in mussels. These results collectively show that novel microbial communities exist between cultured and in-stream freshwater mussels and that their gut bacterial communities may be independent from the influence of environmental factors.</p>	
<p>Afternoon Break 3:10 – 3:25 p.m.</p>	
<p>3:25 – 3:45 p.m.</p>	<p>Disease Risk Analysis – Applications for the Management of Fresh Water Mussels Tiffany Wolf, Phil Miller, Alex Primus & Dominic Travis</p>
<p>Disease risk analysis (DRA) is a structured, evidence-based process that helps decision making in the face of uncertainty by characterizing the potential impact of infectious and non-infectious diseases on ecosystems, wildlife, domestic animals and people. Since 1992 the Conservation Planning Specialist Group (CPSG, previously the Conservation Breeding Specialist Group) of the IUCN Species Survival Commission (IUCN-SSC) has been facilitating collaboration between experts in zoo and wildlife veterinary medicine, disease ecology and population management to develop a set of tools for analysis of wildlife disease risks. In 2010, CPSG, in partnership with three other IUCN-SSC specialist groups (Wildlife Health, Reintroduction and Invasive Species), undertook a global needs analysis survey in this area. The results demonstrated that wildlife disease concerns are global, broad in scope and involve a wide diversity of people from multiple disciplines. The resulting <i>Manual of Procedures for Wildlife Disease Risk Analysis</i> built upon a large body of work on <i>disease risk analysis</i>, in particular that of the World Organisation for Animal Health (OIE), and extends this to biodiversity conservation. The University of Minnesota College of Veterinary Medicine’s Risk Analysis Unit (https://www.cahfs.umn.edu/risk-analysis-service) brings together expertise and resources across the University of Minnesota to develop science and risk-based tools to meet new challenges in the areas of animal health, food safety and wildlife. In partnership, CPSG and UMN have developed a mini workshop series aimed at helping conservationists, animal resource managers and industry professionals integrate science and policy in order to frame, characterize and manage health risks using these international standards. Participation in this workshop will help managers and policy makers: transition health policy problems into a scientific risk-based framework; organize and communicate with diverse stakeholder working groups; design a structured approach to identifying and prioritizing disease hazards; develop a first iteration conceptual risk assessment model.</p>	

3:50 – 4:05 p.m.	DO THE INFECTIOUS DISEASES OF DREISSENIDS REPRESENT A THREAT TO NORTH AMERICAN UNIONID POPULATIONS? Daniel P. Molloy
<p>The spread of dreissenids (zebra and quagga mussels) across North America has had a significant negative impact on unionid populations. High densities of dreissenids settling on their shells often lead to the death of unionids, resulting in severe population declines. But is there any evidence that these invasive mussels might also represent an additional threat to unionids due the parasites they carry? This presentation will examine what parasites dreissenids have been documented to carry in North America and in their native Eurasia and what evidence there is that these parasites are crossing over to infect unionids on these continents.</p>	
4:10 – 4:25 p.m.	INVASIVE SNAILS AND THEIR POTENTIAL TO SERVE AS HOSTS FOR PARASITES IN THE MIDWEST Gregory J Sandland
<p>Abstract: Invasive species have important impacts on the ecology and economics of regions across the world. Snails are an important group of aquatic invaders because they have the potential to transmit parasites that are problematic for both wildlife and people. As part of this presentation I will 1) highlight some of the key invasive snails found in freshwater systems throughout the Midwest and 2) outline their potential to serve as hosts for flatworm parasites in the region.</p>	
4:30 – 4:45 p.m.	FACTORS RELATED TO GROWTH INHIBITION IN JUVENILE MUSSELS EXPOSED TO AMBIENT STREAM CONDITIONS. Wendell R. Haag, Jacob Culp, Monte McGregor, Steven Price & Lesley Sneed
<p>In a previous study, caged juvenile mussels survived but did not grow in streams that have previously lost their mussel fauna; growth was as much as two orders of magnitude higher in streams that continue to support mussels. Growth inhibition was associated with agricultural contaminants (pesticides, nitrates) in some streams but not in others, and no potential causal factor was common to all streams. We measured growth of hatchery-reared juvenile mussels exposed to ambient conditions in 10 streams, five that previously showed growth inhibition and five that did not. We examined other potential causal factors and other measures of performance associated with growth inhibition. Diatom assemblages showed a geographical pattern but no relationship to growth. Microbe assemblages showed no pattern related to geography or growth; these results do not support differences in food resources as explanations for low growth. C and N stable isotope signatures showed that mussels with growth inhibition assimilated available food resources to a lesser extent than mussels with normal growth. Mussels that showed growth inhibition had a distinctive metabolomic profile in which citric acid cycle activity and amino acid metabolism were greatly reduced, but fatty acid oxidation was elevated; these traits distinguished those individuals reliably from individuals with normal growth. This profile likely reflects a starvation response in which lipids are catabolized to provide energy in the absence of energy normally produced via the citric acid cycle. The factors responsible for this metabolomic response and failure to assimilate available food resources are unknown at this time.</p>	

4:50 – 5:05 p.m.	1999 OHIO RIVER MOLLUSK KILL ASSESSMENT: THE GASTROPOD STORY. Janet L. Clayton & Patricia A. Morrison
<p>An extensive mollusk kill occurred on the Ohio River in 1999 resulting from a toxic release of hexavalent chromium and the treatment thereof with a molluscicide compound. The kill originated in the Belleville Pool below Marietta, OH. The kill assessment was conducted by Ecological Specialists and consisted of semi-quantitative transects and excavated quadrats at five known mussel beds below the alleged discharge. Impacts were observed nearly 30 miles downstream. While the primary objective was to assess the unionid kill, the quadrat excavations provided an opportunity to also assess the impact to the snails. The dominant snail observed was <i>Pleurocera canaliculata</i> with three additional species observed. It was believed that 100% mortality of unionids occurred at the most upstream assessed bed, and total mortality at the five beds was estimated at 870,000 individuals; approximately 11 million gastropods died at those sites alone. Losses in between and beyond those sites are immeasurable. Both unionid and gastropod recovery has been monitored over the past 18 years. Snails were much slower to rebound and whereas the population was primarily made up of <i>P. canaliculata</i> prior to the kill it is now composed of <i>P. canaliculata</i> and <i>Lithasia verrucosa</i>. An explosion of <i>Birgella subglobosus</i> occurred in 2007 but has only barely persisted since. Episodic settlement and growth of zebra mussels continue to threaten the full recovery of both the unionids and gastropods.</p>	
5:10 – 5:30 p.m.	FRESHWATER MUSSEL DIE-OFFS: INSIGHTS FROM A COMPILATION OF KNOWN CASES Jordan Richard
<p>In 1986 a symposium was conducted to address the emerging threat of unexplained mass die-offs of freshwater mussels in the United States. Although the symposium brought significant attention to the topic at the time, publications and records of die-offs in subsequent years are rare. As part of an effort to focus the investigation of the ongoing die-off of freshwater mussels in the Clinch River, TN, we attempted to review data and conclusions from all documented die-off events. We issued calls for information through both the unio listserv and the FMCS mailing list. Including the data from the 1986 workshop, our compilation presently contains 48 cases of varying magnitude from seven countries. The cases include a number of events with known causes, but also include many mass die-offs that went entirely unexplained. Based on significant spatial and temporal biases in the cases reported, we believe the list is largely incomplete, and are actively seeking additional responses. Here, we discuss some of the most significant and well-documented cases, and share evidence of insights derived from commonalities in the case reports. Among these, we find evidence that: 1) Die-offs driven by various mechanisms leave unique “signatures” that can guide investigations based on simple, inexpensive analysis of an in-progress die-off; 2) Die-offs often persist over multiple years; and 3) Many of the populations affected by die-offs have never recovered.</p>	
Poster Session & Social 7:00-9:00	

WEDNESDAY MARCH 14, 2018

<p>7:30 – 9:30 a.m.</p>	<p align="center">CONTINENTAL BREAKFAST</p>	
<p>8:00 – 11:30 a.m. 12:00 – 4:30 p.m.</p>	<p>LABORATORY SESSION: pp.19 <i>8:00AM BUS LEAVES GROUP 1(PINK HEELSPLITTERS- red dot on nametag)</i> <i>12:00PM BUS LEAVES AT GROUP 2(GREEN FLOATERS-green dot on nametag)</i></p>	<p>Expert Leads: Eric Leis, Corey Puzach, Katie Bockrath, Susan Knowles, Andrew McElwain</p>
<p>8:00 – 11:30 a.m. 1:00 – 4:30 p.m.</p>	<p>RISK CHARACTERIZATION SESSION: pp. 21 <i>8:00AM- GROUP 2 (GREEN FLOATERS-green dot on nametag)</i> <i>1:00PM- GROUP 3 (YELLOW SANDSHELL- yellow dot on nametag)</i></p>	<p>Expert Leads: Tiffany Wolf, Alex Primus and Phil Miller Facilitator: Megan Bradley</p>
<p>8:00 – 11:30 a.m. 1:00 – 4:30 p.m.</p>	<p>DIE-OFFS AND KILLS SESSION: pp. 20 <i>8:00AM - GROUP 3 (YELLOW SANDSHELL- yellow dot on nametag)</i> <i>1:00PM- GROUP 1(PINK HEELSPLITTERS- RED DOT ON NAMETAG)</i></p>	<p>Expert Leads: Heidi Dunn, Dr. Nick Phelps & Jordan Richards Facilitator: Diane Waller</p>
<p align="center">Social & Jam Session 7:00-9:00</p>		

THURSDAY MARCH 15, 2018

7:30 – 9:30 a.m.	CONTINENTAL BREAKFAST	
8:00 – 11:30 a.m.	<p>LABORATORY SESSION: pp.19 <i>8:00AM BUS LEAVES GROUP 3 (YELLOW SANDSHELL- yellow dot on nametag)</i></p>	<div style="border: 1px solid black; padding: 5px;"> <p>Expert Leads: Eric Leis, Corey Puzach, Katie Bockrath, Susan Knowles, Andrew McElwain</p> </div>
8:00 – 11:30 a.m.	<p>RISK CHARACTERIZATION SESSION: pp. 21 <i>8:00AM- GROUP 1(PINK HEELSPLITTERS- red dot on nametag)</i></p>	<div style="border: 1px solid black; padding: 5px;"> <p>Expert Leads: Tiffany Wolf, Alex Primus and Phil Miller Facilitator: Megan Bradley</p> </div>
8:00 – 11:30 a.m.	<p>DIE-OFFS AND KILLS SESSION: pp. 20 <i>8:00AM- GROUP 2 (GREEN FLOATERS-green dot on nametag)</i></p>	<div style="border: 1px solid black; padding: 5px;"> <p>Expert Leads: Heidi Dunn, Dr. Nick Phelps & Jordan Richards Facilitator: Diane Waller</p> </div>
1:00 – 5:00 p.m.	FIELD TRIP TO GENOA NATIONAL FISH HATCHERY	

Laboratory Session Outline

Eric Leis, Corey Puzach, Katie Bockrath, Susan Knowles, Andrew McElwain

- I. Introduction-Microscope use
- II. Mussel Anatomy
 - A) Overview of mussel organs and tissues
 - 1. Mantle
 - 2. Gills and glochidia
 - 3. Labial palps
 - 4. Integument of the visceral mass
 - 5. Foot
 - 6. Digestive gland
 - 7. Stomach
 - 8. Crystalline style sac
 - 9. Intestine
 - 10. Gonads
 - 11. Nephridium
 - 12. Heart
 - 13. Hemocytes and hemolymph
 - 14. Ganglia and nerves
 - 15. Shell damage and shell deformities
 - B) Sampling specific tissues for histology
 - 1. Sampling from live animals
 - 2. Sampling from fixed animals
 - 3. Sampling from infected/lesioned tissue
 - 4. Grossing/trimming tissue samples
 - C) Optimizing images
 - 1. White balance
 - 2. Brightness
 - 3. Scale bar
 - 4. Optimizing photographs using Adobe Photoshop
 - 5. Making figures or plates with Adobe Illustrator
- III. Histological samples
 - A) Proper fixation and storage of mussels for histology
 - 1. Prop shell open with wooden dowels
 - 2. Cut adductor muscles
 - 3. Remove adductor and mantle from one side of the shell
 - 4. Long term storage in ethanol
 - 5. Comparing fixation/preservation methods (formalin, ethanol, freezing)
 - B) Proper fixation of snails and small clams
- IV. Bacteriology
- V. Virology
- VI. Parasitology
 - A) How to perform a necropsy
 - 1. Open shell
 - B) Cursory inspection of mantle cavity
 - 1. Remove and inspect specific tissues
 - i. Mantle
 - 2. Gill
 - 3. Integument of visceral mass
 - 4. Foot
 - 5. Anterior visceral mass
 - 6. Posterior visceral mass
 - 7. Dorsal visceral mass

Mollusk Die-off Session

Dr. Ryan Carnegie (VIMS) – Mortality events across a continuum: lessons from marine bivalve disease outbreaks

Dr. Nick Phelps (UMN) – Standardizing and simplifying fish kill reporting

Heidi Dunn (Ecoanalysts) – Investigation and monetary values of fish and freshwater mollusk kills
Effective sampling design and techniques for assessment of mussel populations

Jordan Richard (USFWS – Virginia) – Review of a case study: Clinch River mussel die-off

Disease Risk Analysis – Applications for the Management of Freshwater Mussels

Tiffany Wolf, Phil Miller, Alex Primus, & Dominic Travis

Background and Resources

The following brief overview and resources are included as a user-friendly guide to outline the process of Disease Risk Analysis (DRA) and define common terminology. For a more thorough introduction to the DRA, participants are referred to the IUCN's *Manual of Procedures for Wildlife Disease Risk Analysis*.

Overview of the Process

The DRA consists of five general steps, with risk communication with stakeholders being a key component throughout (Figure 1). Risk analysis refers to the overall process; distinct from Risk Assessment, a stage of the process. DRA starts with problem description (the process of describing and justifying the problem or question) and follows with Hazard Identification, Risk Assessment, Risk Management, and Implementation and Review. Again, Risk Communication should occur frequently throughout the process and engage technical experts, key stakeholders, and policy makers to maximize the quality of the analysis and enhance the likelihood of implementation of resulting recommendations. Problem Description is the first step of the process and critical to the success of the DRA. In this step, participants should clearly define the context of the problem, the specific question to be addressed, the goal and scope of the DRA, assumptions and limitations, and the acceptable level of risk. Hazard Identification involves the identification of all possible hazards, both infectious and non-infectious. Criteria should be developed related to the defined problem to aid in prioritizing the list of hazards, consequences, both direct and indirect, of each of the hazards should be considered to identify which should be further evaluated. The Risk Assessment stage of the DRA is a more formal approach to assessing 1) the likelihood of release (introduction) of a hazard into a system, 2) the likelihood of the species of interest being exposed to the hazard following release and 3) the consequences of exposure. This may be accomplished through qualitative, semi-quantitative, or quantitative methods. Risk Management entails a review of potential risk reduction or management strategies for mitigation that may be employed along with their potential outcomes. This forms the basis for DRA recommendations. Finally, the DRA process concludes with development of a plan for the process and timeline for monitoring, evaluation and review of the proposed risk management actions.




Figure 1: Overview of the Disease Risk Analysis process taken from the IUCN Manual of Procedures for Wildlife Disease Risk Analysis

Glossary of Terms- Select terminology and definitions taken from IUCN *Manual of Procedures for Wildlife Disease Risk Analysis*.

acceptable risk	A level of risk that is so small in terms of likelihood of occurrence or consequences that, in comparison with the expected benefits, stakeholders are willing to accept it
consequence assessment	The process of describing the relationship between specified exposures to a hazard and the consequences of those exposures. A causal process must exist by which exposures produce adverse health or environmental consequences, which may in turn lead to socioeconomic consequences and consequences for conservation. The consequence assessment describes the consequences of a given exposure and estimates the probability of them occurring
disease	Any impairment of the normal structural or physiological state of a living organism resulting from its physiological response to a <i>hazard</i>
disease risk analysis	The application of <i>risk analysis</i> to identify diseases that may enter a specified animal population to identify the likelihood of such introductions, assess their consequences and identify measures that may be applied to mitigate either the likelihood of introduction or the magnitude of consequences
endemic	A disease or parasite the prevalence of which does not exhibit wide fluctuations through time in a defined location. The term ‘enzootic’ is sometimes applied when referring to non-human populations
epidemic	A sudden, rapid spread or increase in the prevalence or intensity of a parasite or disease. An epidemic is often the result of a change in circumstances that favor parasite transmission such as a rapid increase in host population density or the introduction of a new parasite. Having an established baseline is essential for detecting epidemics. The term ‘epizootic’ is sometimes applied when referring to non-human populations

exposure assessment	The process of describing the biological pathway(s) necessary for exposure of animals and humans in a particular environment to the hazards (in this case the pathogenic agents) released from a given risk source, and estimating the probability of the exposure(s) occurring, either qualitatively or quantitatively
hazard	A biological, chemical or physical agent in, or a condition of, an animal or animal product with the potential to cause an adverse health effect. See also <i>disease</i>
hazard identification	The process of identifying the pathogenic or hazardous agents that could potentially be introduced into a specified animal population or environment by the activity being considered
host	Any animal that is capable of harboring a parasite, regardless of whether it plays a role in the further transmission of the parasite
incidence	The number of new health events (infection, disease, etc.) experienced by a given population over a specific period of time. (cf. prevalence, the total number, new and old, in a given population in a specified time period)
incubation period	The time that elapses between infection with a parasite and the onset of disease
infection	The entry and development or multiplication of a parasite in the body of a host, where it may or may not cause disease
infectious period	Period during which the infected individual is able to transmit the infection
infestation	Subsistence of a macroparasite on the external surface of a host regardless of whether the infestation results in disease
latent period	The period when an individual is infected but not yet capable of transmitting the infection

macroparasites	Parasites that in general do not multiply within their hosts but instead produce transmission stages (eggs and larvae) that pass into the external environment (e.g. the parasitic helminths (worms) and arthropods).
model	In the context of DRA, a graphical or computational representation of an actual system used to predict disease dynamics and impacts, and the effect of management interventions on those dynamics and impacts
monitoring	The intermittent performance and analysis of routine measurements and observations, aimed at detecting changes in the environment or health status of a population
parasite	An agent that lives on or within a host and that survives at the expense of the host regardless of whether a disease state follows. This definition includes both microparasites (e.g. bacteria, viruses) and macroparasites (e.g. helminths, arthropods)
pathogen	Any disease-causing <i>parasite</i>
prevalence	The proportion of the host population with infection, disease or antibody presence, often expressed as a percentage. A measure of how widespread an infection, disease or exposure to an infectious agent is at a point in time
qualitative risk assessment	An assessment in which the outputs on the likelihood of the outcome or the magnitude of the consequences are expressed in qualitative terms such as high, medium, low or negligible
Quantitative risk assessment	An assessment in which the outputs of the risk assessment are expressed numerically
release assessment	The process of describing the biological pathway(s) necessary for a particular activity to ‘release’ (that is, introduce) hazards into a particular environment or ecosystem, and estimating the probability, either qualitatively or quantitatively, of that complete process occurring

reservoir	Any animate (humans, animals, insects, etc.) or inanimate object (plant, soil, feces, etc.) or any combination of these serving as a habitat of a parasite that reproduces itself in such a way as to be transmitted to a susceptible host
risk	The likelihood of the occurrence and the likely magnitude of the consequences (biological, economic, etc. as defined by a specific risk analysis question) of an adverse event or effect to animal or human health
risk analysis	The process composed of <i>problem description, hazard identification, risk assessment, risk management</i> and <i>risk communication</i>
risk assessment	The evaluation of the likelihood and the consequences of entry, establishment or spread of a pathogenic agent within a specified animal population or environment
risk communication	The interactive exchange of information and opinions throughout the <i>risk analysis</i> process concerning risk, risk-related factors and risk perceptions among risk assessors, risk managers, risk communicators, the general public and other interested parties
risk management	The process of identifying, selecting and implementing measures that can be applied to reduce the level of risk
surveillance	The systematic ongoing collection, collation and analysis of information related to animal health and the timely dissemination of information to those who need to know so that action can be taken
uncertainty	The lack of precise knowledge of the input values that is due to measurement error or to lack of knowledge of the steps required, and the pathways from hazard to risk, when building the scenario being assessed
vector	An insect or any living carrier that transports an infectious agent from an infected individual to a susceptible individual or its food or immediate surroundings. The organism may or may not pass through a development cycle within the vector
zoonosis	A disease naturally transmitted between humans and other vertebrate species

POSTER SESSION ABSTRACTS, TUESDAY MARCH 13, 2018- 7:00PM

<p>Die-off Section Poster 1</p>	<p>BIG DARBY CREEK, OHIO, 2016 MUSSEL DIE-OFF <u>A.M. Sasson</u>¹, G.T. Watters², A. Boose³, D. Symonds², J. Gordon⁴, C. Byrne². ¹<i>The Nature Conservancy, Dublin, OH 43017</i>; ²<i>The Ohio State University, Columbus, OH 43212</i>; ³<i>Columbus and Franklin County Metro Parks, Westerville, OH 43081</i>; ⁴<i>Lawhon & Associates, Columbus, OH 43212</i>.</p> <p>In late 2016 and early 2017, a large die-off of mussels across all species was observed in Ohio’s Big Darby Creek, a tributary in the Scioto River basin. Big Darby Creek is a State and National Scenic River due to its outstanding fish and mussel diversity, whose 144,000 ha (555 mi²) watershed is dominated by agricultural land use. The die-off was first observed in October 2016, with lingering effects continuing into early 2017. An unusually high number of distressed and fresh dead mussels were observed throughout ~93 km (~58 mi) of Big Darby Creek. Based on about 200 person-hours of survey and collection efforts, the number of fresh dead shells observed and collected at many sites was estimated at often one to two orders of magnitude greater than has been known to be observed by well-practiced observers familiar with mussels in this watershed. Of the 43 recorded species and 35+ known to be presently extant in the watershed, it appeared that all species of freshwater mussels were affected in the Big Darby Creek die-off, from the most common (e.g., spike, kidneyshell, fatmucket, plain pocketbook) to federally endangered (northern riffleshell, clubshell). Mussel deaths appeared to occur over a period of weeks to months, ending by spring 2017. There were no known observations of fish mortality, and no cause was determined through state and federal agency water and tissue sampling. Six sites were surveyed on Little Darby Creek although a die-off was not observed in this tributary.</p>
<p>Die-off Section Poster 2</p>	<p>THE DIE-OFFS OF FRESHWATER PEARL MUSSEL (<i>MARGARITIFERA MARGARITIFERA</i>) IN SWEDEN; AN OVERVIEW OF THE CURRENT SITUATION AND POSSIBLE CAUSES. <u>Niklas Wengström</u>^{1,2}, Håkan Söderberg³, Johan Höjesjö¹. ¹<i>University of Gothenburg, Department of Biological and Environmental Sciences, Box 463, SE-405 30 Gothenburg, Sweden</i>; ²<i>Swedish Anglers Association, Sjölyckan 6, 416 55 Gothenburg, Sweden</i>; ³<i>The county administrative board of Västernorrland, Pumpbacksgatan 19 871 86 Härnösand</i>.</p> <p>The freshwater pearl mussel <i>Margaritifera margaritifera</i> is an endangered species in Sweden with more than 600 known populations distributed in 16 counties. Approximately one third of these populations can be considered viable and healthy as they show sign of recent recruitment of juveniles (<50 mm). The problem with recruitment failure in Sweden is associated with turbidity and sedimentation of fine substrates and other alterations in their habitat. Almost every running water has been used historically for lodging and/or mills and since the beginning of the 20th century hydropower plants has been a problem. In the late 1960’s acidification of streams and lakes became one of Sweden’s biggest environmental problem and in 1977 a liming program started to mitigate the effects of acidification in surface waters. Many of these problems have been solved or are about to get solved. Since the 1980’s Sweden have measured freshwater pearl mussel populations with quantitative methods and we use juvenile recruitment as one factor indicating a healthy environment. Recently, there have been reports of freshwater pearl mussel die-offs in five different counties resulting in population decreases between 22 – 100 %. The underlying reason for this sudden death in many systems is not clear but we believe that it is due to both biotic and abiotic factors. We have therefore started a small-scale monitoring project aiming to detect possible factors and we will in this presentation provide a map visualizing the current knowledge of where freshwater pearl mussel die-offs have been found in Sweden.</p>

<p>Die-off Section Poster 3</p>	<p>EU LIFE IP PROJECT FRESHABIT: CONSERVATION OF THE TWO MAIN REMAINING <i>MARGARATIFERA MARGARATIFERA</i> POPULATIONS OF SOUTHERN FINLAND <u>Jouni Taskinen</u>¹, Jukka Pakkala², Eero Mäenpää², Anu Suonpää³, Juha-Pekka Vähä³, Esko Vuorinen⁴ and Panu Oulasvirta⁵. ¹<i>Department of Biological and Environmental Science, University of Jyväskylä, Box 35, FI-40014 JYU, Finland;</i> ²<i>Centre for Economic Development, Transport and the Environment of South Ostrobothnia, Finland;</i> ³<i>Association for Water and Environment of Western Uusimaa, Finland;</i> ⁴<i>Silvestris Nature Surveys Ltd., Finland.</i> ⁵<i>Alleco Ltd., Finland.</i></p> <p>The freshwater pearl mussel (eastern pearl shell, <i>Margaritifera margaritifera</i>, FPM) is endangered throughout the range of its distribution in Europe. There are over 120 rivers inhabiting FPM in Finland, most of them being located in the north. Only two major FPM rivers are found from the southern Finland – River Ähtävänjoki/Esseen and River Mustionjoki/Svartå, flowing to the Baltic Sea. The estimated FPM population size of River Ähtävänjoki has decreased from 50,000 to 500 from between 1987-2016. Reproduction of FPM was evident in River Ähtävänjoki from 1990 to 2006, but not in 2015-16. Steep decline of FPM was evident also in River Mustionjoki, from 3,000 to 1,300 between 2010-2016, accompanied with cessation of glochidia release. I will describe the planned/on-going actions of the EU Life IP project FRESHABIT (2016-2021) to prevent extinction of these last remaining southern Finnish FPM populations: transportation of adult mussels to tank facility to resume glochidia production by using artificial feeding, captive breeding, habitat restoration and fishway construction to restore the natural migration and reproduction of salmonid hosts.</p>
<p>Die-off Section Poster 4</p>	<p>EFFECTIVENESS OF TWO ANTIFUNGAL MEDICATIONS ON FUNGAL INFECTION IN SOUTHERN COMBSHELL, <i>EPIOBLASMA PENITA</i> (CONRAD, 1834) <u>Michael Buntin</u>¹, Todd Fobian¹, Benjamin Beck², and Paul Johnson¹. ¹<i>Alabama Department of Conservation, Alabama Aquatic Biodiversity Center, Marion, AL, 36756;</i> ²<i>USDA-Agricultural Research Service, Auburn, AL 36832.</i></p> <p>Southern Combshell is a federally endangered unionid endemic to the Mobile River Basin in Alabama and Mississippi. Propagation of this species has been taking place at the Alabama Aquatic Biodiversity Center since 2010. In 2017, five female Southern Combshells taken from the Buttahatchee River, Lowndes County, MS became infested with fungal hyphae after two individuals had their glochidia extracted via water-filled syringe. The animals were initially grouped by species population in separate aerated 1.5 L covered containers stored in an 8°C communal incubator. Once the infection was observed, the Southern Combshells were quarantined in a separate incubator. Three individuals died before any treatment could be administered. The remaining two mussels were treated independently with Pimafix™ and amphotericin B. The amount of visible hyphae on the mantle or foot was reduced on both animals after multiple days of treatment but neither eliminated the infection entirely. Both treated Southern Combshells eventually died after the visible hyphae returned to pretreatment levels. No fungal hyphae were observed on any other mussel within the original brood stock incubator. The fungi were later identified by Dr. Cova Arias, using ITS-2 sequencing, as <i>Didymella bellidis</i> and <i>Cladosporium sphaerospermum</i>. While not a controlled experiment, the inability of these medications to eliminate the fungal infestation in these mussels emphasizes the need for sterile procedures and isolation of brood stock to prevent the spreading of potentially lethal infections among individuals and populations.</p>
<p>Die-off Section Poster 5</p>	<p>DISENTANGLING THE DIFFERENT ROLE OF PARASITES AND INVASIVE SPECIES IN THE DECLINE OF NATIVE MUSSELS <u>Nicoletta Riccardi</u>¹, Maria Urbanska², Wojciech Andrzejewski², Angela Boggero¹, Caterina Fioroni¹, Jouni Taskinen³. ¹<i>Institute of Ecosystem Study, National Research Council, Italy;</i> ²<i>Poznan University of Life Sciences, Poland;</i> ³<i>University of Jyväskylä, Finland.</i></p> <p>Parasites have important negative consequences for freshwater mussels, but in the past they have been regarded as non-lethal and without any drastic impact upon mussel populations. Nonetheless, increased parasite densities impair mussel physiological conditions, which causes an increased mortality rate in mussels exposed to other stressors. A recently reported stressor is the spread of non-indigenous species (NIS), that can introduce new parasites and/or alter the dynamics of endemic parasites. To explore the interaction between biological invasions and growing parasites presence we profited of the sharp decline of the dominant native mussel (<i>Unio elongatulus</i>) in Lake Maggiore (Northern Italy), after the arrival of <i>Corbicula fluminea</i> and <i>Sinanodonta woodiana</i>. This study addressed whether: 1) the native species decline was determined by NIS co-introduction of own parasites; 2) introduced species triggered endemic parasite emergence; 3) habitat characteristics affect the interacting impacts of NIS and parasites. To answer these questions we evaluated:</p> <ol style="list-style-type: none"> 1) the occurrence of parasites (alien/endemic) and the frequency of shared parasites in invasive and native bivalve species; 2) the prevalence and intensity of parasitic diseases in native species: <ol style="list-style-type: none"> a. before and after NIS arrival (task 2); b. in different habitat types and at different depths (task 3)

Poster 6	<p>MACROSCOPICAL AND MICROSCOPICAL FINDINGS FROM STUDIES OF ONGOING DECLINES IN SWEDISH FRESHWATER PEARL MUSSEL (<i>MARGARITIFERA MARGARETIFERA</i>) POPULATIONS ANDERS ALFJORDEN¹, Håkan Söderberg², Tomas Troschke³. ¹National Veterinary Institute, Sweden (SVA), S-751 89 Uppsala; ²County administrative board of Västernorrland, S-871 86 Härnös; ³County administrative board of, Gävleborg, S-801 70 Gävle.</p>
	<p>Several cases of increased mortality in freshwater pearl mussels (<i>Margaritifera margaritifera</i> L. 1758) have been detected during surveys of freshwater creeks and rivers during the last years (2011-2017) in Sweden. No etiological cause of these wild stock mortalities has been concluded yet. In some cases, samples from live animals have been collected for necropsy and histological investigations, during field surveys. During recent field surveys (2016-2017), freshwater mussels have been transported alive for further investigations to the National Veterinary Institute in Uppsala. These mussels have been compared with mussels collected the same day from healthy population where no signs of mortalities have been reported. Ocular screening of cross sections of these animals was investigated regarding pathological findings. Freshwater pearl mussels from affected population showed clear pathological changes compared to reference mussels both regarding macroscopical and microscopical appearances. The mussels from healthy population had firm and thick bodies, well developed gonadal organs and digestive glands filled with dark green digested feed content. In comparison the mussel from the affected population was in general much thinner with more relaxed bodies and had a lower body weight. Furthermore, the affected, lethargic mussels showed low or no presence of gonad cells (egg cells) and the digestive gland was pale with watery and transparent content. The microscopical investigation confirmed this pattern showing changes in the affected mussels compared to the reference population. The mussels from the declining populations had empty gonad follicles, enlarged swollen epithelial cells in digestive glands, where cells showed no presence of granular or vesicular content. Sometimes cellular infiltration was seen in the connective tissue surrounding these digestive glands. In the solid external body parts where connective and muscle fibers dominated, an increased cellular infiltration was observed in the emaciated mussels.</p>
Poster 7	<p>NON-DESTRUCTIVE DIAGNOSIS OF TREMATODE PARASITISM IN FRESHWATER MUSSELS JOUNI TASKINEN, Jocelyn Mah Choo, Markus Tulolsela & Benjami Kumpalalnen</p>
	<p>Freshwater mussels (Unionoida) are declining throughout the world, possibly signaling poor health status of mussels. Digenean trematodes of the family Bucephalidae and Gorgoderidae (e.g. <i>Bucephalus</i>, <i>Rhipidocotyle</i>, <i>Proisorhynchoides</i> and <i>Phyllodistomum</i> spp.) infect unionid clams, producing thousands of cecarial larvae and causing sterility of mussel host. Microscopical examination of mussels for trematodes requires destructive dissection of mussel tissues. Therefore, an alternative, non-destructive method would be useful as endangered mussels can only rarely be sacrificed for a parasitological survey. We studied if cercariae shedding could reliably reveal <i>Rhipidocotyle fennica</i> infection in the mussel <i>Anodonta anatina</i>. In the 1st experiment, individual mussels were kept in 2 L aquaria for 1 h, after which the water was checked for cercariae and the tissues examined microscopically. 26% of infections were found in ambient water temperature of 17 C, but 81% when temperature was increased to 21 C. Second experiment was conducted at field by incubating mussels in three time points during the cercariae shedding season, and by using water temperature 3 C higher than the ambient lake water. 63 % of infections were detected during the seasonal peak shedding period (mid-August) while on average 44% of infected individual were detected outside the optimal season (late July and late August). Thus, cercaria method could serve as a non-invasive way to investigate trematode infections in freshwater mussels.</p>
Poster 8	<p>COMMUNITY-BASED MONITORING OF FRESHWATER MUSSELS AS A RESEARCH AND CONSERVATION PRACTICE Celeste A. Searles Mazzacano. CASM Environmental, LLC, Portland, OR 97206.</p>
	<p>Oregon is home to three genera of native freshwater mussels: <i>Margaritifera falcata</i> (Western Pearlshell), <i>Gonidea angulata</i> (Western Ridged Mussel), and several clades of <i>Anodonta</i> (Floater), including the Oregon/Western Floater clade (<i>A. oregonensis/kennerlyi</i>) and California/Winged Floater clade (<i>A. californiensis/nuttalliana</i>). Oregon's freshwater mussels are known to be declining, but they are also more poorly characterized than many other aquatic fauna, and detailed or long-term studies are lacking for most water bodies. With no federal protections in place for western mussels, funding mechanisms and legal obligations that help drive research and conservation are lacking. In addition, many practitioners do not know whether mussel populations are present in the waters they manage, and/or they lack the resources to characterize known or suspected populations. Community-based monitoring done by volunteers trained and overseen by content experts can be a powerful and cost-effective tool to improve freshwater mussel research, conservation, and education. This poster will describe best practices in community-based freshwater mussel monitoring and use case studies in Oregon to illustrate the benefits and challenges of these projects.</p>

<p>Poster 9</p>	<p>TOXICITY EVALUATION OF AN EFFLUENT ENTERING THE DEEP FORK RIVER, OK, USA JAMES KUNZ¹, Jeff Steevens¹, Ning Wang¹, Suzanne Dunn² and David Martinez². ¹US Geological Survey, Columbia, MO 65201; ²US Fish and Wildlife Service, Tulsa, OK.</p> <p>Effluents from a local manufacturing company were suspected as the cause of two mussel kills in 2005 and in 2011 in the Deep Fork River (DFR), OK. Previous studies reported elevated potassium in water contaminated by the effluent. The objectives of this study were to (1) assess the potential toxicity of the effluent to a unionid mussel (<i>Lampsilis siliquoidea</i>) and two commonly tested species (cladoceran <i>Ceriodaphnia dubia</i>; fathead minnow <i>Pimephales promelas</i>) in 7-day effluent tests; (2) evaluate the relative sensitivities of the three species to potassium in 7-day toxicity tests; and (3) determine the influence of water hardness on the acute toxicity of potassium in reconstituted waters. Preliminary results indicated that the effluent-contaminated water contained 50 times more potassium than upstream DFR water and was highly toxic to mussels and cladocerans. The mussel was more sensitive than the cladoceran and fathead minnow in all tests. The acute toxicity of potassium to the mussel and cladoceran did not substantially change in four reconstituted waters mimicking water quality characteristics in the DFR with a broad hardness range of 35 to 300 mg/L as CaCO₃, and importantly, the 50% effect concentrations for mussels were below the potassium concentrations measured in the effluent in this and other studies.</p>
<p>Poster 10</p>	<p>THE VALVE MOVEMENT RESPONSE OF <i>UNIO TUMIDUS</i> AND <i>SINANODONTA WOODIANA</i> EXPOSED TO NITRATE, SULFATE CHLORATE AND PHOSPHATE COMPOUNDS Joanna Chmyst¹, <u>Maria Urbańska</u>². ¹Department of Ecology & Environmental Protection; ²Institute of Zoology, Poznań University of Life Sciences, ul. Wojska Polskiego 28, 60-637 Poznań, Poland.</p> <p>For last years bivalves have presented one of the most widely used organisms in monitoring water quality establishing the biology early warning system (BEWS). We compared the sensitivity of European native <i>Unio tumidus</i> and alien, <i>Sinanodonta woodiana</i> bivalves. <i>U. tumidus</i> has been used as a water safety monitor for more than 8 million people in Poland. <i>S. woodiana</i> is one of the species that has recently been reported as being invasive worldwide. KNO₃ (50 mg l⁻¹), K₂SO₄ (250 mg l⁻¹), NH₄Cl (250 mg l⁻¹) and KH₂PO₄ (0,75 mg l⁻¹) were tested in the continuous recirculation system. For each experiment 8 individuals of mussels were placed into the system. Both species react in similar way for each exposure to pollution taking into account both mean daily shell opening level and mean daily activity time. For the chloride and phosphate pollution there is no reaction in this measure for both species. The greatest reactions are for nitrate and sulfate exposure. While the mean daily shell opening level declines twice for <i>U. tumidus</i>, this leap is 12.5 times for <i>S. woodiana</i>. However, in absolute values this fall is smaller by 48% and 23%, respectively, so the detection may be easier in the former case. The mean daily activity time drops is by 57% and as much as 88% for <i>U. tumidus</i> and <i>S. woodiana</i>, respectively.</p>
<p>Poster 11</p>	<p>IS IT POSSIBLE TO HIDE FROM <i>DREISSENA POLYMORPHA</i>? EXAMPLE OF NATIVE AND INVASIVE SPECIES OF UNIONIDAE FOULING BY ZEBRA MUSSELS <u>Maria Urbanska</u>¹, Wojciech Andrzejewski¹ and Henryk Gierszal². ¹Institute of Zoology, Poznan University of Life Sciences, ul. Wojska Polskiego 28, 60-637 Poznań, Poland; ²Department of Applied Computer Science, Adam Mickiewicz University, Umultowska 85, 61 -614 Poznań, Poland.</p> <p><i>Unionidae</i> live partly buried in the sediments, usually with the posterior part of their shells exposed to the water column to enable filtration. This exposed part of the shell can therefore serve as a substratum for zebra mussels, which limit the host's ability to feed, move, and breed. We analyzed images of shells of 2 unionid species: <i>Anodonta anatina</i> and alien <i>Sinanodonta woodiana</i>. 40 specimens of both species fouled or not by <i>D. polymorpha</i> were randomly collected. The recorded data comprised approximate area of the whole shell, the exposed part, and of the outer parts of <i>D. polymorpha</i>, protruding from the unionid shell. Our results show that these 2 species significantly differed in shell surface available to zebra mussels. <i>A. anatina</i> was consistently more heavily fouled than <i>S. woodiana</i> and had a greater surface area of the shell exposed in the water column. On average only 25.5% of the shell of <i>S. woodiana</i> was exposed above the sediments, compared to 45.4% of the shell of <i>A. anatina</i>. Thus <i>S. woodiana</i> is strongly buried in the mud, which gives it better protection from the negative impact of zebra mussels and makes the Chinese species more competitive than the native <i>A. anatina</i>.</p>

<p>Poster 12</p>	<p>EFFECT OF TEMPERATURE ON SURVIVAL, GROWTH AND DEVELOPMENT OF MARGARITIFERA MARGARITIFERA LARVAE DURING EARLY PARASITISM IN SALMO SALAR GILLS. <u>Castrillo, P. A.¹, Varela C.², Ronza, P.¹, Outeiro A.², Bermúdez R.¹, Quiroga M.I.¹, Ondina P.².</u> ¹ <i>Department of Anatomy, Animal Breeding and Veterinary Clinical Sciences;</i> ² <i>Department of Zoology, Genetics and Physical Anthropology. Universidade de Santiago de Compostela, Facultad de Veterinaria, 27002, Lugo, España.</i></p> <p><i>Margaritifera margaritifera</i> is a freshwater mussel categorized as Critically Endangered and protected by European conservation policies. Still, the propagation programs are constrained by its life cycle, characterized by a long-term and specific parasitic stage of the larvae (glochidia) on fish gills. It is known that temperature influences the metabolism and immune response of bivalves and fish, nevertheless, the effect of this factor on the interaction between fish and glochidia has not been investigated. The purpose of this study was to determine the effect of temperature on <i>M. margaritifera</i> larvae during the early parasitic stages on <i>Salmo salar</i> gills. Two fish groups were exposed to glochidia and maintained at 9 and 17 °C, respectively, for 30 days. Parr were euthanized and second left holobranches were assessed by stereomicroscopy. The number and the maximal diameter of glochidia between both groups were compared by Wilcoxon and ANOVA test, respectively. Moreover, remaining holobranches were processed for histological examination. In fish exposed at 17°C, larvae displayed higher number ($P<0.05$), bigger size ($P<0.01$) and more advanced organogenesis than those at 9°C. These results suggest that temperature influences survival, growth and development rates during early parasitic stages of <i>M. margaritifera</i>. Moreover, combination of stereomicroscopical and histological studies represent a good tool to study the host-parasite interaction, although further studies are needed to understand the subjacent mechanisms and the consequences in older stages of this freshwater pearl mussel. This work has been funded by the <i>MarMaCul</i> project from the Fundación Biodiversidad and by a predoctoral contract of Xunta de Galicia.</p>
<p>Poster 13</p>	<p>DEVELOPING KEY SKILLS IN FRESHWATER MUSSEL CONSERVATION AT THE NATIONAL CONSERVATION TRAINING CENTER (USFWS) <u>Matthew Patterson.</u> <i>National Conservation Training Center, United States Fish and Wildlife Service, 698 Conservation Way, Shepherdstown, WV 25443</i></p> <p>The plight of native freshwater mussels in North America and around the world has led to significant increases in the number of natural resource professionals spending all or a portion of their time on mussel conservation issues. Within the United States Fish and Wildlife Service alone, Ecological Services biologists are being asked to review project proposals and new listings, hatchery biologists are being asked to propagate mussels for recovery of threatened and endangered species, and national wildlife refuge biologists are being asked to survey mussel populations on their property. Unfortunately, many of these biologists have never received formal training in the biology, ecology, life history, or identification of freshwater mussels. A similar lack of formal training can be found at other federal, state, and non-governmental organizations. To help meet this training need, the National Conservation Training Center (NCTC) in Shepherdstown, West Virginia developed a freshwater mussel curriculum that currently includes three courses, Conservation Biology of Freshwater Mussels, Freshwater Mussel Propagation for Restoration, and Freshwater Mussel Identification. Since starting the curriculum in 2012, the NCTC has trained over 250 biologists in freshwater mussel biology, ecology, life-history, field sampling, propagation and culture, and identification in an effort to improve conservation efforts on the ground and stem the tide of population declines. This poster presentation will provide an overview of the mussel curriculum, a break-down of students trained by state and agency, as well as information on upcoming courses.</p>
<p>Poster 14</p>	<p>THE MUSSELS OF LOLO CREEK <u>Doug Nemeth, Frank Mullins, Mike Murray, Chris Griffith.</u> <i>Idaho Fish and Wildlife Conservation Office, Orofino, Idaho.</i></p> <p>Freshwater mussels are one of the most imperiled groups of animals in the world, so when a habitat restoration project in Lolo Creek, Clearwater River Basin, Idaho was going to result in dewatering a large number of western pearlshell mussels (<i>Margaritifera falcata</i>), agencies agreed to cooperatively translocate the mussels prior to dewatering. This provided an opportunity to investigate mussel quantification techniques and the demographic characteristics of Lolo Cr. mussels. Objectives were to estimate the numbers of mussels present using two different techniques and compare those estimates to the actual census (number of mussels translocated), determine the proportion of mussels on the surface to those buried and their lengths, and determine the recovery rate of PIT-tagged, translocated mussels. Floating downstream and counting all mussels within the prescribed area was more efficient than using quadrats and a systematic random sampling design. The number and proportion of mussels on the surface in Lolo Cr. was strongly influenced by water temperature. Mussels may have segregated by length through the substrate column. Disturbance and removal of mussels resulted in additional mussels recruiting to those areas. Recovery rate of PIT-tagged mussels was about 50% at 4°C, but may be higher at warmer temperatures when more mussels are on the surface.</p>

<p>Poster 15</p>	<p>PHYSIOLOGICAL PROFILE OF NATIVE UNIONID MUSSELS IN RESPONSE TO ZEBRA MUSSEL INVASION <u>Vince L. Butitta, Alyssa Ginther & Emily H. Stanley. Center for Limnology, University of Wisconsin-Madison, Madison, WI 53706.</u></p> <p>One of the fastest growing threats to North America’s native freshwater mussel is the invasion of Eurasian Zebra mussel (<i>Dreissena polymorpha</i>). Since the introduction and rapid spreading of invasive Dreissenid mussels throughout North America, native Unionid species have experienced dramatic declines—with many reports of local extirpation, but some situations of possible coexistence. Despite well-documented declines in native mussel populations to Dreissenid invasion, the mechanisms through which Unionids are locally extirpated is not well understood. Zebra mussel populations began exploding in the Mendota Lake (Dane Co., WI) in 2016, they have since quickly colonized the three lakes downstream. We collected individuals from two common Unionid species (<i>Lampsilis siliquoidea</i> and <i>Pyganodon grandis</i>) across a Zebra mussel infestation gradient. Hemolymph samples were extracted for metabolomic analysis, soft tissues were dried, weighed, and assessed for glycogen content. Glycogen has long been used to quantify general poor body condition in mussels, but the powerful new tool of metabolomics has recently been shown to be extremely useful in understanding how mussels physiologically respond to stressful handling, hypoxia, food limitation, and certain toxicants. A metabolomic analysis of Unionids suffering from a zebra mussel invasion may provide a more detailed look into the mechanism through which Unionid populations collapse following zebra mussel invasions and could offer insight into possible coexistence.</p>
<p>Poster 16</p>	<p>DEVELOPMENT OF WEB-BASED TAXONOMIC KEY TO IDENTIFY FRESHWATER GLOCHIDIA. <u>Heidi Dunn¹, Kathy Koch², Nate Jacobson², Doug Bradley², Chris Cieciek², Jonathon Black³, Doug Dixon³. ¹EcoAnalysts, Inc, O’Fallon, MO 63366; ²LimnoTech, Ann Arbor, MI 48108; ³Electric Power Research Institute, New Salem, MA 01355.</u></p> <p>Taxonomists traditionally use a variety of hardcopy guides to identify freshwater unionid mollusk larvae (glochidia). These guides may not include contemporary identification characteristics and/or current nomenclature, may lack high resolution photographs or images, and may be out of print or difficult to obtain. Further, new morphological information observed at propagation facilities may not be published. With Electric Power Research Institute (EPRI) funding, a conceptual web-based lucid key has been designed to aid researchers and professionals with identification of freshwater unionid mollusk glochidia. The web-based platform is modeled on other taxonomic keys (freshwater and marine larval fish) developed by the EPRI and LimnoTech team and is intended to provide a single, open resource that supports the compilation and dissemination of taxonomic information, allows for the incorporation of new and high-quality information and imagery, and facilitates improved identification. The project is presently adding species-level information and images to the conceptual key framework to improve the database and support the accurate identification of freshwater glochidia. Current activities also include collaboration with malacology experts to identify additional data and image resources to further enhance the database, and broaden the exposure and access of glochidia identification among scientists.</p>
<p>Poster 17</p>	<p>PARASITISM IN UNIONID MUSSELS: A THREAT FOR SPECIES AT RISK <u>Joseph P. Carney & Andrea Strawson. Department of Biology, Lakehead University, Thunder Bay, ON, P7B 5E1, Canada.</u></p> <p>Freshwater mussels can host a variety of helminth parasites, including digenean sporocysts. Host sterility, can result from sporocyst infection. The purpose of this study was to assess the helminth parasite fauna of freshwater mussels from the Lake Winnipeg drainage. A total of 1021 freshwater mussels (Bivalvia: Unionidae) representing 12 species collected from 35 rivers in Manitoba, Saskatchewan and North Dakota was examined for helminth parasites. Five helminth species infected a total of 10 mussel species. Adult <i>Aspidogaster conchicola</i> infected 5 mussel species, and adult <i>Cotylogaster occidentalis</i> infected 9 mussel species with no apparent effect to the hosts. Metacercariae, a larval digenean, parasitized 6 mussel species, with no apparent effect on the host. Five mussel species were infected with 2 types of digenean sporocyst. A sporocyst releasing rhopalocercous cercariae infected 3 mussel species with no apparent effect on mussel reproduction. A second sporocyst, releasing furcocercous cercariae, infected 5 mussel species and was responsible for either reducing mussel reproductive capacity, or completely sterilized the mussel. Subsequent histological study of <i>Pyganodon grandis</i> demonstrated the absence of gonad tissue associated with heavy infection by this second type of sporocyst. Decreasing reproduction, either partially or completely, both lowers individual fitness, and decreases the effective breeding size of the population. These parasite induced pathologies need to be considered when conservation efforts are directed toward mussel species at risk.</p>

Poster 18	<p>PROPAGATION AT VIRGINIA FISHERIES AND AQUATIC WILDLIFE CENTER BEN DAVIS¹, Amy Maynard², Rachel Mair¹, Brian Watson³, Michael Odom¹, Bryce Maynard¹. ¹<i>U.S. Fish and Wildlife Service, Charles City, VA 23030</i>; ²<i>Conservation Management Institute, Virginia Polytechnic & State University, Charles City, VA 23030</i>; ³<i>Virginia Department of Game and Inland Fisheries, Forest, VA 24551</i>.</p>
	<p>Virginia Fisheries and Aquatic Wildlife Center (VFAWC) is a cooperative freshwater mussel propagation facility located at Harrison Lake National Fish Hatchery. Since VFAWC's founding in 2007, the US Fish and Wildlife Service and the Virginia Department of Game and Inland Fisheries have jointly propagated 12 species of Atlantic slope freshwater mussels, and have released over 247,000 individuals back into the wild. During the 2017 propagation season, VFAWC produced over 1,152,000 juvenile mussels of 8 species of special concern, including the federally-endangered <i>Pleurobema collina</i> (James spiny mussel) and the state-threatened <i>Lasmigona subviridis</i> (green floater). Of the mussels produced, approximately 228,000 were distributed to other hatcheries or researchers. A total of 45,421 mussels of 7 species were released during 2017 as part of population augmentation efforts. Future work will include mitigation projects utilizing both conventional and in-vitro propagation methods.</p>
Poster 19	<p>REINTRODUCTION OF EPIOBLASMA AUREOLA (BIVALVIA: UNIONIDAE) IN THE UPPER CLINCH RIVER BASIN, TAZEWELL COUNTY, VIRGINIA. Sarah L. Colletti, Tim W. Lane, Joseph F. Ferraro & Tiffany C. Leach. <i>Aquatic Wildlife Conservation Center, Virginia Department of Game and Inland Fisheries, Marion, Virginia 24354</i>.</p>
	<p>The critically endangered Golden Riffleshell mussel <i>Epioblasma aureola</i> (Jones and Neves 2010) is an Upper Tennessee River Basin (UTRB) endemic restricted to a single reproducing population in the Upper Clinch River. Previous population augmentation efforts using host fish were met with limited success and wild brood stock have become increasingly challenging to locate. In 2016, VDGIF biologists recovered three gravid females from Indian Creek, VA and transferred their glochidia to KDFWR's Center for Mollusk Conservation, Franklin, KY. There, using <i>in vitro</i> methods, 12,000 juveniles were successfully produced. At ~3 months of age, 1,200 remaining juveniles were split between 3 culture facilities with variable survival but were eventually transferred for final grow-out to VDGIF's Aquatic Wildlife Conservation Center (AWCC), Marion, VA in 2017. To initiate pre-release monitoring, 5 sites across the upper Clinch and Indian Creek were chosen in 2017 for silo deployment. Two silos with five mussels each were placed at all sites and monitored monthly for growth and survival. Survival was 100% across all sites, with mussels exhibiting variable growth among sites. The UTRB Mussel Recovery Group selected three out of the five sites for release, one being the native brood stock location in Indian Creek and two others in the main stem Clinch River. Pit-tagged <i>E. aureola</i> (N=100) were released in a 2.5 m² grid at each site and color-coded with Hallprint© tags to identify maternal pedigree. An additional 431 individuals were free-released across the chosen sites, for a total of 731 sub-adults released through this augmentation effort. Silos remain at the five sites and those 50 individuals will continue to be monitored. Mark-recapture surveys are planned to observe the growth and survival of the pit-tagged individuals. Individuals from the 2013 (N=4) and 2016 (N=252) cohorts, and a new 2017 (N=39) cohort, remain in captivity and are being molecularly analyzed by Virginia Tech University. This data is intended to inform potential avenues for the creation of an ARK population and eventual captive breeding program at AWCC. Future efforts to expand the species' range beyond the Clinch River watershed are planned. This accomplishment and future efforts are only possible because of extensive interagency cooperation and collaboration.</p>

<p>Poster 20</p>	<p>PESTS, PREDATORS, PARASITES, AND RELATED PROBLEMS IN THE CULTURE OF FRESHWATER MUSSELS <u>Morgan Kern</u>¹, Louise Lavictoire¹, Megan Bradley², Nathan Eckert², Beth Glidewell³ & Chris Barnhart¹. ¹<i>Department of Biology, Missouri State University, Springfield, MO 65897</i>; ²<i>Genoa National Hatchery, Genoa WI 54632</i>; ³<i>Confederated Tribes of the Umatilla Indian Reservation, Pendleton, OR, 97801</i>.</p> <p>A wide variety of organisms impact the captive culture of freshwater mussels. We provide an overview of some of these organisms, their effects, and mitigation methods. Parthenogenic midge larvae (<i>Paratanytarsus</i>) immobilize juveniles and foul screens by case building. Midge can be controlled with entomopathic bacteria products such as Gnatrol[®]. Suspension feeding Cladocera and Ostracoda similar in size to juvenile mussels may bloom in culture. If removal by filtration is impractical, these can be reduced by decanting as they swim above the substrate. Colonial ciliates and rotifers can foul surfaces including juveniles as well as competing for suspended food. Routine control involves cleaning culture vessels, screens and substrate. Fouling organisms can be removed from juveniles by brief drying. Predators on early juveniles include oligochaete annelids (<i>Chaetogaster</i>) and rhabdocoel flatworms (<i>Macrostomum</i>, <i>Microstomum</i>). Rhabdocoels can be removed from host fish with formalin. Water molds, probably mainly <i>Saprolegnia</i>, are occasionally observed in dead or moribund juveniles and adults. Sterilizing larval trematodes (Bucephalidae) occur sporadically in Unionidae and may affect from 0-25% of wild adult mussels. Completion of the trematode life cycle requires host fish, so that reinfection is unlikely but possible in pond-based culture operations. Diseases that commonly affect host fish carrying mussels include <i>Flavobacterium columnaris</i>, which responds to Kanamycin sulfate and NaCl. The ciliate <i>Ichthiophthirius</i> can be controlled with NaCl, which excysting juvenile mussels can tolerate briefly up to 4.5 g/L. Praziquantel can control <i>Monogenea</i> in host fish without harm to juveniles.</p>
<p>Poster 21</p>	<p>A COMPARISON OF SUBSTRATE PREFERENCES FOR NATIVE AND INVASIVE MUSSEL POPULATIONS IN THE KISHWAUKEE RIVER <u>Sophie C. McComb</u> & <u>Madeline M. McCormick</u>. <i>Students, Sycamore High School, Sycamore, IL 60178</i>.</p> <p>Freshwater mussels (family Unionidae) are particularly important for the preservation of local freshwater aquatic habitats and environments. Of increasing priority is examining the relationship between the growing population of non-native <i>C. fluminea</i> and the survival of the species in aquatic environments, as well as the discovery of any substrate preferences. Many of the articles reviewed indicate that the scientific community studying freshwater mussels has paid great attention to the latter, but very few studies focus on a relationship between non-native <i>C. fluminea</i> and native mussel populations. Two sites were surveyed within the East Branch of the South Branch of the Kishwaukee River sub-watershed. A total of 64 randomly selected quadrats were surveyed for substrate composition, native mussel species, and non-native mussel populations. Significant correlations were found between both native and nonnative populations to the percentage of sand in the substrate, as well as a strong positive correlation between <i>C. fluminea</i> populations and the presence of native species. Important in the consideration of these results is the late period of survey in the freshwater mussel activity season, and the relatively small sample size of the study. The results, with consideration to the limitations of the study, suggest the dominance and strong adaptive ability of <i>Corbicula</i> (Sousa, et al. 2009). This also presents a need for further examination of the relationships between native and non-native species, and the containment of invasive population growth.</p>
<p>Poster 22</p>	<p>CONSERVATION OF MARGARITIFERA MARGARITIFERA IN SPAIN: AN OVERVIEW OF THE MARGALULLA PROJECT Varela, C.¹, CASTRILLO, P.A.², Outeiro, A.¹, Amaro, R.¹, San Miguel, E.¹, Castro, J.¹, Ondina, P.¹. ¹<i>Dpto. de Zoología, Genética y Antropología Física</i>; ²<i>Dpto de Anatomía Patológica y Ciencias Clínicas Veterinarias. Facultad de Veterinaria. Universidad de Santiago de Compostela. 27002 Lugo. España</i>.</p> <p>The naiad <i>M. margaritifera</i> is an endangered species throughout its holartic distribution area. Galician river basins (NW Spain, on the southwestern limit of its European range) are a core area that houses 80% of Iberian populations. However, Galician rivers have suffered an increased number of pressures during the last century, which have significantly decreased their quality and ecological function. As a consequence, most of the mussel populations are now fragmented, with low abundance and little or no recruitment currently. The European Life funds provides financing for environmental protection and nature conservation projects with the aim of implementing and developing the EU's environmental policies. MargalUlla was conceived as a long-term species conservation project, dealing both with captive breeding and habitat restoration in the Ulla basin. Accordingly, a breeding program has been launched in 2013 in the rearing facility "O Veral" (the only one on the Iberian Peninsula) with the aim of obtaining juveniles to reinforce populations.</p> <p>In this work we present the results obtained in different experiences carried out over four years. The publication of this kind of data is essential to understand conservation measures on wild populations and to improve the efficiency of applied methods of rearing. This work has been funded by the <i>MarMaCul</i> project from the Fundación Biodiversidad (Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente). The experience was carried out at Veral rearing facility (Consellería de Medio Ambiente, Xunta de Galicia).</p>

Poster 23	<p>ASSESSING THE IMPACT OF CONTAMINANTS OF EMERGING CONCERN ON THE HEALTH AND REPRODUCITON OF FRESHWATER MUSSELS: TOOLS TO INFORM RESOURCE MANAGEMENT</p> <p>Mandy Annis¹ & <u>Daelyn A. Woolnough</u>². ¹U.S. Fish and Wildlife Service, Michigan Ecological Services Office, East Lansing MI 48823. ²Department of Biology and Institute for Great Lakes Research, Central Michigan University, Mt. Pleasant MI, 48859.</p>
	<p>Contaminants are considered a factor in freshwater mussel decline with increasing interest in the impacts of contaminants of emerging concern (CEC). CECs including pharmaceuticals, personnel care products, manufacturing by-products, and new agricultural chemicals are found in complex mixes in all landscapes including mussel habitats that support species of concern, yet the impacts of CECs are unknown. Research often does not accurately reflect real world exposure of mussels with complex lifecycles and long lives. Further, research regarding sub-lethal, populationlevel impacts of CECs found at non-lethal levels are limited. Understanding and non-lethally assessing the impacts of CECs on declining populations is vital to species conservation. With funding through the Great Lakes Restoration Initiative, the US Fish and Wildlife Service and Central Michigan University, among others, have partnered to develop tools which begin to address these knowledge gaps. Studies include mussel and host fish surveys, stream side exposures of mussels to naturally experienced CEC gradients, and laboratory exposures of mussels and host fish to ecologically relevant chemical mixes through brooding, host fish interaction, and transformation. Biological, behavioral, and contaminants data have been collected and will be evaluated to assess mussel health in relation to CECs. Non-lethal tool development includes adapted biological sampling, and population and hazard assessment modeling. These studies will provide information regarding potential population-level impacts of CEC mixtures enabling managers to select optimal conservation and management practices in order to avoid further population declines, future listing of imperiled species, and loss of trust resources.</p>

Bibliography

Marine Bivalve Pathology:

- Allam, B., and Raftos, D. 2015. Immune responses to infectious diseases in bivalves. *Journal of Invertebrate Pathology* 131: 121–136.
- Carella, F., Feist, S.W., Bignell, J.P., and De Vico, G. 2015. Comparative pathology in bivalves: Aetiological agents and disease processes. *Journal of Invertebrate Pathology* **131**: 107–120. doi:10.1016/j.jip.2015.07.012.
- Carnegie, R.B., Arzul, I., and Bushek, D. 2016. Managing marine mollusc diseases in the context of regional and international commerce: policy issues and emerging concerns. *Philosophical Transactions of the Royal Society B: Biological Sciences* **371**(1689): 20150215. doi:10.1098/rstb.2015.0215.
- Jones-Lepp, T.L., Taguchi, V., Sovocool, W., Betowski, D., DeArmond, P., Schumacher, B., Winnik, W., McMillin, R., and Armstrong, C. 2018. Novel contaminants identified in fish kills in the Red River watershed, 2011–2013. *Environ Toxicol Chem* **37**(2): 336–344. doi:10.1002/etc.3989.
- Metzger, M.J., Reinisch, C., Sherry, J., and Goff, S.P. (n.d.). Horizontal Transmission of Clonal Cancer Cells Causes Leukemia in Soft-Shell Clams. *Cell* **161**(2): 255–263. doi:10.1016/j.cell.2015.02.042.
- Powell, E.N., and Hofmann, E.E. 2015. Models of marine molluscan diseases: Trends and challenges. *Journal of Invertebrate Pathology* **131**: 212–225. doi:10.1016/j.jip.2015.07.017.
- Zannella, C., Mosca, F., Mariani, F., Franci, G., Folliero, V., Galdiero, M., Tiscar, P.G., and Galdiero, M. 2017. Microbial Diseases of Bivalve Mollusks: Infections, Immunology and Antimicrobial Defense. *Marine Drugs* 15(6): 182.

Marine Bivalve Virology:

- Arzul, I., Corbeil, S., Morga, B., and Renault, T. (n.d.). Viruses infecting marine molluscs. *Journal of Invertebrate Pathology*.
- Shi, M., Lin, X.-D., Tian, J.-H., Chen, L.-J., Chen, X., Li, C.-X., Qin, X.-C., Li, J., Cao, J.-P., Eden, J.-S., Buchmann, J., Wang, W., Xu, J., Holmes, E.C., and Zhang, Y.-Z. 2016. Redefining the invertebrate RNA virosphere. *Nature* 540(7634): 539–543.

Gastropods:

- Michelson, E. H. 1970. *Aspidogaster conchicola* from freshwater gastropods in the United States. *J. Parasitol.*, 56: 709-712.

Immunology:

- Evariste, L., Auffret, M., Audonnet, S., Geffard, A., David, E., Brousseau, P., Fournier, M., and Betoulle, S. 2016. Functional features of hemocyte subpopulations of the invasive mollusk species *Dreissena polymorpha*. *Fish & Shellfish Immunology* 56: 144–154. doi:10.1016/j.fsi.2016.06.054.
- Gélinas, M., Fortier, M., Lajeunesse, A., Fournier, M., Gagnon, C., and Gagné, F. 2013. Energy status and immune system alterations in *Elliptio complanata* after ingestion of cyanobacteria *Anabaena flos-aquae*. *Ecotoxicology* 22(3): 457–468. doi:10.1007/s10646-012-1039-4.
- Hine, P 1999. The interrelationship of bivalve hemocytes. *Fish and Shellfish Immunology* 9:367-385
- Roch, P. 1999. Defense mechanisms and disease prevention in farmed marine invertebrates. *Aquaculture* 172(1): 125–145. doi:10.1016/S0044-8486(98)00439-6.

Freshwater Mollusk Health:

- Archambault J.M., Gregory Cope W. & Kwak T.J. (2013) Burrowing, byssus, and biomarkers: Behavioral and physiological indicators of sublethal thermal stress in freshwater mussels (Unionidae). *Marine and Freshwater Behaviour and Physiology* **46**, 229–250.
- Bartsch M.R., Bartsch L.A., Richardson W.B., Vallazza J.M. & Moraska Lafrancois B. (2017) Effects of food resources on the fatty acid composition, growth and survival of freshwater mussels. *Plos One* **12**, e0173419.
- Berg D.J., Haag W.R., Guttman S.I. & Sickel J.B. (1995) Mantle biopsy: a technique for nondestructive tissue-sampling of freshwater mussels. *Journal of the North American Benthological Society* **14**, 577–581.
- Fritts A.K., Peterson J.T., Hazelton P.D. & Bringolf R.B. (2015a) Evaluation of methods for assessing physiological biomarkers of stress in freshwater mussels. *Canadian Journal of Fisheries and Aquatic Sciences* **72**, 1450–1459.

Bibliography

- Fritts A.K., Peterson J.T., Wisniewski J.M. & Bringolf R.B. (2015b) Nonlethal assessment of freshwater mussel physiological response to changes in environmental factors. *Canadian Journal of Fisheries and Aquatic Sciences* **72**, 1460–1468.
- Gustafson L.L., Stoskopf M.K., Bogan A.E., Showers W., Kwak T.J., Hanlon S., *et al.* (2005a) Evaluation of a nonlethal technique for hemolymph collection in *Elliptio complanata*, a freshwater bivalve (Mollusca: Unionidae). *Diseases of Aquatic Organisms* **65**, 159–165.
- Gustafson L.L., Stoskopf M.K., Showers W., Cope G., Eads C., Linnehan R., *et al.* (2005b) Reference ranges for hemolymph chemistries from *Elliptio complanata* of North Carolina. *Diseases of Aquatic Organisms* **65**, 167–176.
- Hannan K.D., Jeffrey J.D., Hasler C.T. & Suski C.D. (2016a) Physiological effects of short- and long-term exposure to elevated carbon dioxide on a freshwater mussel, *Fusconaia flava*. *Canadian Journal of Fisheries and Aquatic Sciences* **53**, 1538–1546.
- Hannan K.D., Jeffrey J.D., Hasler C.T. & Suski C.D. (2016b) Physiological responses of three species of unionid mussels to intermittent exposure to elevated carbon dioxide. *Conservation Physiology* **4**, 1–13.
- Hannan K.D., Jeffrey J.D., Hasler C.T. & Suski C.D. (2016c) The response of two species of unionid mussels to extended exposure to elevated carbon dioxide. *Comparative Biochemistry and Physiology-Part A: Molecular and Integrative Physiology* **201**, 173–181.

Freshwater Mollusk Disease:

- Butros, J. 1948. A tumor in a fresh-water mussel. *Cancer Research* **8**: 270–271.
- Beecher, C. E. 1884. Some abnormal and pathologic forms of fresh-water shells from the vicinity of Albany, New York. *New York State Museum Annual Report*. **36**: 51–55 with five plates.
- Carella, F., Villari, G., Maio, N., and De Vico, G. 2016. Disease and Disorders of Freshwater Unionid Mussels: A Brief Overview of Recent Studies. *Frontiers in Physiology* **7**.
- Grizzle, J. M. and Brunner, C. J. 2009. Infectious diseases of freshwater mussels and other freshwater bivalve mollusks. *Reviews in Fisheries Science [Rev. Fish. Sci.]* **17**(4):425-467.
- Pauley, G. B. 1967. A tumorlike growth on the foot of a freshwater mussel (*Anodonta californiensis*). *Journal of the Fisheries Research Board of Canada* **24**: 679–682.
- Pauley, G. B. 1968. A disease of the freshwater mussel, *Margaritifera margaritifera*. *Journal of Invertebrate Pathology* **12**: 321–328.
- Pekkarinen, M. 1993. Reproduction and condition of unionid mussels in the Vantaa River, South Finland. *Archiv für Hydrobiologie* **127**: 357–375.
- Roper, D. S., and C. W. Hickey. 1994. Population structure, shell morphology, age and condition of the freshwater mussel *Hyridella menziesi* (Unionacea: Hyriidae) from seven lake and river sites in the Waikato River system. *Hydrobiologia* **284**: 205–217.
- Strayer, D. L. 2008. A widespread morphological deformity in freshwater mussels from New York. *Northeastern Naturalist* **15**: 149–151.

Freshwater Bivalve Parasitology:

- Alves, P. V., F. M. Vieira, C. P. Santos, T. Scholz, and J. L. Luque. 2015. A checklist of the Aspidogastrea (Platyhelminthes: Trematoda) of the world. *Zootaxa* **3918**: 339–396.
- Anderson, R. V., and D. J. Holm. 1987. *Chaetogaster limnaei* (Oligochaeta: Naididae) infesting unionid mollusks (Pelecypoda: Corbiculidae) in Pool 19, Mississippi River. *Journal of Freshwater Ecology* **4**: 61–64.
- Antipa, G. A. and E. B. Small. 1971. The occurrence of thigmotrichous ciliated protozoa inhabiting the mantle cavity of unionid molluscs of Illinois. *Trans. Am. Microsc. Soc.*, **90**: 463-472.
- Aristanov, E. 1992. The role of *Dreissena polymorpha* Pallas in the life cycle of *Bucephalus polymorphus* Baer 1827. *Uzb. Biol. Zh.*, 1992: 75-76 (in Russian).
- Baker, F. C. 1901. Some interesting molluscan monstrosities. *Transactions of the Academy of Science of St. Louis*. **11**: 143–146 with one plate.
- Bakker, K. E. and C. Davids. 1973. Notes on the life history of *Aspidogaster conchicola* Baer, 1826 (Trematoda: Aspidogastridae). *J. Helminthol.*, **47**: 269-276.

Bibliography

- Baturo, B. 1977. *Bucephalus polymorphus* Baer, 1827 and *Rhipidocotyle illense* (Ziegler, 1883) (Trematoda, Bucephalidae): Morphology and biology of developmental stages. *Acta Parasitol. Pol.*, 24: 203-220 + Plates I-IV.
- Bradbury, P. C. 1994. Parasitic protozoa of molluscs and crustacea. In: *Parasitic Protozoa* (Vol. 8, 2nd ed.), pp. 139-263. (J. P. Kreier, Ed.). San Diego, California: Academic Press.
- Burlakova, L. E., Karatayev, A. Y., and Molloy, D. P. 1998. Field and laboratory studies of zebra mussel (*Dreissena polymorpha*) infection by the ciliate *Conchophthirus acuminatus* in the Republic of Belarus. *J. Invertebr. Pathol.* 71:251-257
- Carney, J.P. 2015. Aspidobothrean Parasites of Freshwater Mussels (Bivalvia: Unionidae) from the Saskatchewan–Nelson River Drainage in Manitoba, Canada and North Dakota, United States. *Comparative Parasitology* 82(1): 9–16.
- Chernysheva, N. B., and K. J. Purasjoki. 1991. A redescription of *Paraergasilus rylovi* Markevich, 1937 (Copepoda, Ergasilidae). *Systematic Parasitology* 20: 165–171.
- Clark, H. W., and C. B. Wilson. 1912. The mussel fauna of the Maumee River. Bureau of Fisheries Document No. 757. 72 pp.
- Coker, R. E., A. F. Shira, H. W. Clark, and A. D. Howard. 1921. The Fisheries Biological Station at Fairport, Iowa. U.S. Bureau of Fisheries Document 895: 1–12.
- Conn, D. B. and D. A. Conn. 1995. Experimental infection of zebra mussels *Dreissena polymorpha* (Mollusca: Bivalvia) by metacercariae of *Echinoparyphium* sp. (Platyhelminthes: Trematoda). *J. Parasitol.*, 81: 304-305.
- Conn, D. B., A. Ricciardi, M. N. Babapulle, K. A. Klein, and D. A. Rosen. 1996. *Chaetogaster limnaei* (Annelida: Oligochaeta) as a parasite of the zebra mussel *Dreissena polymorpha*, and quagga mussel *Dreissena bugensis* (Mollusca: Bivalvia). *Parasitol. Res.*, 82: 1-7.
- Cribb, T. H., R. A. Bray, P. D. Olson, and D. T. J. Littlewood. 2003. Life cycle evolution in the Digenea: a new perspective from phylogeny. *Advances in Parasitology* 54: 197–254.
- Curry, M. G. 1977. Delaware leeches (Annelida: Hirudinea: Glossiphoniidae): New state records and new molluscan host record for *Placobdella montifera* Moore. *The Wasmann Journal of Biology* 35: 65–67.
- Curry, M. G., B. Everitt, and M. F. Vidrine. 1981. Haptobenthos on shells of living freshwater clams in Louisiana. *The Wasmann Journal of Biology* 39: 56–62.
- Davids, C. and M. H. S. Kraak. 1993. Trematode parasites of the zebra mussel (*Dreissena polymorpha*). In: *Zebra Mussels: Biology, Impacts, and Control*, pp. 749-759. (T. F. Nalepa and D. W. Schloesser, Eds.). Boca Raton: Lewis Publishers.
- Edwards, D.D., and Vidrine, M.F. 2006. Host specificity among unionicola spp. (acari: unionicolidae) parasitizing freshwater mussels. *Journal of Parasitology* 92(5): 977–983.
- Edwards, D. D., and M. F. Vidrine. 2013. Mites of freshwater mollusks. Malcolm F. Vidrine, Eunice Louisiana, 336 p.
- Fenchel, T. 1965. Ciliates from Scandinavian molluscs. *Ophelia*, 2: 71-174.
- Forsyth, D. J., and I. D. McCallum. 1978. *Xenochironomus canterburyensis* (Diptera: Chironomidae), a commensal of *Hyridella menziesi* (Lamellibranchia) in Lake Taupo; features of pre-adult life history. *New Zealand Journal of Zoology* 5: 759–800.
- Fuller, S. L. H. 1974. Clams and Mussels (Mollusca: Bivalvia). In *Pollution Ecology of Freshwater Invertebrates*, C. W. Hart, Jr., and S. H. Fuller, (eds.). Academic Press, New York. p. 215–273.
- Gangloff, M.M., Lenertz, K.K., and Feminella, J.W. 2008. Parasitic mite and trematode abundance are associated with reduced reproductive output and physiological condition of freshwater mussels. *Hydrobiologia* 610(1): 25–31
- Gentner, H. W., and S. H. Hopkins. 1966. Changes in the trematode fauna of clams in the Little Brazos River, Texas. *Journal of Parasitology* 52: 458–461.
- Gordon, M. J., B. K. Swan, and C. G. Paterson. 1978. *Baeoctenus bicolor* (Diptera: Chironomidae) parasitic in unionid bivalve mollusks, and notes on other chironomid-bivalve associations. *Journal of the Fisheries Research Board of Canada* 35: 154–157.
- Graczyk TK, Thompson RCA, Fayer R, Adams P, Morgan UM, Lewis EJ (1999c) *Giardia duodenalis* of genotype A recovered from clams in the Chesapeake Bay subestuary, Rhode River. *Am J Trop Med Hyg* 61:526–529

Bibliography

- Graczyk TK, Fayer R, Lewis EJ, Higgins JA, Jenkins MC, Thompson RCA, Xiao L, Adams P, Morgan UM, Lal AA (2000) *Cryptosporidium parvum* oocysts and *Giardia duodenalis* cysts in molluscan shellfish. *Acta Parasitol* 45:148
- Graczyk T., Conn, D., Marcogliese, D., Graczyk, H., and Lafontaine, Y. de. 2003. Accumulation of human waterborne parasites by zebra mussels (*Dreissena polymorpha*) and Asian freshwater clams (*Corbicula fluminea*). *Parasitology Research* 89(2): 107–112. doi:10.1007/s00436-002-0729-x.
- Hendrix, S. S., M. F. Vidrine, and R. H. Hartenstine. 1985. A list of records of freshwater aspidogastriids (Trematoda) and their hosts in North America. *Proc. Helminthol. Soc. Wash.*, 52: 289-296.
- Jokela, J., Taskinen, J., Mutikainen, P., and Kopp, K. 2005. Virulence of parasites in hosts under environmental stress: Experiments with anoxia and starvation. *Oikos* 108:156-164.
- Jokela, J., Uotila, L., and Taskinen, J. 1993. Effect of the castrating trematode parasite *Rhipidocotyle fennica* on energy allocation of fresh-water clam *Anodonta piscinalis*. *Functional Ecology [Funct. Ecol.]* 7:332-338.
- Karatayev, A. Y., Burlakova, L. E., Molloy, D. P., and Mastitsky, S. E. 2007. *Dreissena polymorpha* and *Conchophthirus acuminatus*: What can we learn from host-commensal relationships? *J. Shellfish Res.* 26(4):1153-1160.
- Karatayev, A. Y., Mastitsky, S. E., Burlakova, L. E., Karatayev, V. A., Hajduk, M. M., and Conn, D. B. 2012. Exotic Molluscs in the Great Lakes Host Epizootically Important Trematodes. *Journal of Shellfish Research* 31(3): 885–894.
- Karatayev, A. Y., Burlakova, L. E., Molloy, D. P., and Volkova, L. K. 2000. Endosymbionts of *Dreissena polymorpha* (Pallas) in Belarus. *Int. Rev. Hydrobiol.* 85:539-555.
- Karatayev, A. Y., Burlakova, L. E., Molloy, D. P., Volkova, L. K., and Volosyuk, V. V. 2002. Field and laboratory studies of *Ophryoglena* sp. (Ciliata: Ophryoglenidae) infection in zebra mussels, *Dreissena polymorpha* (Bivalvia: Dreissenidae). *J. Invertebr. Pathol.* 79:80-85.
- Karatayev, A. Y., Mastitsky, S. E., Burlakova, L. E., Molloy, D. P., and Vezhnovets, G. G. 2003. Seasonal dynamics of endosymbiotic ciliates and nematodes in *Dreissena polymorpha*. *J. Invertebr. Pathol.* 83:73-82.
- Karatayev, A. Y., Mastitsky, S. E., Molloy, D. P., and Burlakova, L. E. 2003. Patterns of emergence and survival of *Conchophthirus acuminatus* (Ciliophora: Conchophthiridae) from *Dreissena polymorpha* (Bivalvia: Dreissenidae). *J. Shellfish Res.* 22:495-500.
- Karatayev, A. Y., Molloy, D. P., and Burlakova, L. E. 2000. Seasonal dynamics of *Conchophthirus acuminatus* (Ciliophora, Conchophthiridae) infection in *Dreissena polymorpha* and *D. bugensis* (Bivalvia: Dreissenidae). *Eur. J. Protistol.* 36:397-404.
- Kidder, G. W. 1934. Studies on the ciliates from fresh water mussels, I. The structure of the neuromotor system of *Conchophthirus anodontae* Stein, C. Curtus Engl., and C. Magna sp. nov. *Biological Bulletin* 66: 69–90.
- Klemm, D. J. 1976. Leeches (Annelida: Hirudinea) found in North American mollusks. *Malacol. Rev.*, 9: 63-76.
- Laruelle, F., Molloy, D. P., and Roitman, V. A. 2002. Histological analysis of trematodes in *Dreissena polymorpha*: Their location, pathogenicity, and distinguishing morphological characteristics. *J. Parasitol.* 88(5):856-863.
- Laruelle, F., Molloy, D. P., Fokin, S. I., and Ovcharenko, M. A. 1999. Histological analysis of mantle-cavity ciliates in *Dreissena polymorpha*: Their location, symbiotic relationship, and distinguishing morphological characteristics. *J. Shellfish Res.* 18:251-257.
- Levine, T. D., B. K. Lang, and D. J. Berg. 2009. Parasitism of mussel gills by dragonfly nymphs. *The American Midland Naturalist.* 162: 1–6.
- Lopes, L. P. C., D. M. Pimpão, R. M. Takemoto, J. C. O. Malta, and A. M. B. Varella. 2011. *Hysterothylacium* larvae (Nematoda, Anisakidae) in the freshwater mussel *Diplodon suavidicus* (Lea, 1856) (Mollusca, Unioniformes, Hyriidae) in Aripuanã River, Amazon, Brazil. *Journal of Invertebrate Pathology* 106: 357–359.
- Lopes-Lima, M. et al. 2017. Conservation status of freshwater mussels in Europe: state of the art and future challenges. *Biological Reviews* 92: 572–607.
- Lyakhnovich, V. P., A. Y. Karataev, and N. N. Antsipovich. 1983. The effect of water temperature on the rate of infection of *Dreissena polymorpha* with larvae of *Phyllodistomum folium* Olfers in Lake Lukoml'skoe. *Biol. Vnutr. Vod. Inf. Byull.*, 58: 35-38 (in Russian).

Bibliography

- Mastitsky, S.E., and Veres, J.K. 2010. Field evidence for a parasite spillback caused by exotic mollusc *Dreissena polymorpha* in an invaded lake. *Parasitology Research* 106(3): 667–675.
- McElwain, A., Fleming, R., Lajoie, M., Maney, C., Springall, B., and Bullard, S.A. 2016. Pathological Changes Associated with Eggs and Larvae of *Unionicola* sp. (Acari: Unionicolidae) Infecting *Strophitus connasaugaensis* (Bivalvia: Unionidae) from Alabama Creeks. *Journal of Parasitology* 102(1): 75–86.
- Mills, S. C., M. I. Taylor, and J. D. Reynolds. 2005. Benefits and costs to mussels from ejecting bitterling embryos: a test of the evolutionary equilibrium hypothesis. *Animal Behaviour* 70: 31–37.
- Minguez, L., Molloy, D.P., Guérol, F., and Giambérini, L. 2011. Zebra mussel (*Dreissena polymorpha*) parasites: Potentially useful bioindicators of freshwater quality? *Water Research* 45(2): 665–673.
- Molloy, D. P., Giambérini, L., Burlakova, L. E., Karatayev, A. Y., Cryan, J. R., Trajanovski, S. L., and Trajanovska, S. P. 2010. Investigation of the endosymbionts of *Dreissena stankovici* with morphological and molecular confirmation of host species. Pages 227-237 in *The Zebra Mussels in Europe* (Van der Velde, G., Rajagopal, S., and Bij de Vaate, A., eds.). Backhuys Publishers, Leiden.
- Molloy, D. P., Giambérini, L., Morado, J. F., Fokin, S. I., and Laruelle, F. 2001. Characterization of intracytoplasmic prokaryote infections in *Dreissena* sp. (Bivalvia: Dreissenidae). *Dis. Aquat. Org.* 44(3):203-216.
- Molloy, D. P., Giambérini, L., Stokes, N. A., Bureson, E. M., and Ovcharenko, M. A. 2012. *Haplosporidium raabei* n. sp. (Haplosporidia): A parasite of zebra mussels, *Dreissena polymorpha* (Pallas, 1771). *Parasitology* 139(4):463-477.
- Molloy, D. P., Karatayev, A. Y., Burlakova, L. E., Kurandina, D. P., and Laruelle, F. 1997. Natural enemies of zebra mussels: Predators, parasites, and ecological competitors. *Rev. Fisheries Sci.* 5(1):27-97.
- Molloy, D. P., Lynn, D. H., and Giambérini, L. 2005. *Ophryoglena hemophaga* n. sp. (Ciliophora: Ophryoglenidae): A parasite of the digestive gland of zebra mussels *Dreissena polymorpha*. *Dis. Aquat. Org.* 65:237-243.
- Molloy, D. P., V. A. Roitman, and J. D. Shields. 1996. Survey of the parasites of zebra mussels (Bivalvia: Dreissenidae) in northwestern Russia, with comments on records of parasitism in Europe and North America. *J. Helminthol. Soc. Wash.*, 63: 251-256.
- Müller, T., Czarnoleski, M., Labecka, A.M., Cichy, A., Zajac, K., and Dragosz-Kluska, D. 2015. Factors affecting trematode infection rates in freshwater mussels. *Hydrobiologia* 742(1): 59–70.
- Pauley, G. B., and C. D. Becker. 1968. *Aspidogaster conchicola* in mollusks of the Columbia River system with comments on the host's pathological response. *Journal of Parasitology* 54: 917–920.
- Roback, S. S., D. J. Bereza, and M. F. Vidrine. 1979. Description of an *Ablabesmyia* [Diptera: Chironomidae: Tanypodinae] symbiont of unionid fresh-water mussels [Mollusca: Bivalvia: Unionacea], with notes on its biology and zoogeography. *Transactions of the American Entomological Society* 105: 577–620.
- Robinson, J. L., M. J. Wetzel, and J. S. Tiemann. 2017. Some phoretic associations of macroinvertebrates on transplanted federally endangered freshwater mussels. *Northeastern Naturalist* 24: N29–N34.
- Rosen, R., H. Abe, O. Adejumo, K. Ashami, L. Ballou, K. Montgomery, S. Toe, E. Berg, and L. Peng. 2016. Mean intensity and prevalence of *Cotylaspis insignis* (Trematoda: Aspidogastridae) infections in the fat mucket, *Lampsilis radiata luteola* (Bivalvia: Unionidae), from North Elkhorn Creek, a tributary of the Kentucky River, U.S.A. *Comparative Parasitology* 83: 1–5.
- Saarinen, M., and Taskinen, J. 2003. Reduction In The Level Of Infection Of The Bivalve *Anodonta piscinalis* By The Copepod *Paraergasilus rylovi* Using High Temperature And Low Oxygen. *Journal Of Parasitology* 89(6): 1167–1171.
- Saarinen, M., and Taskinen, J. 2004. Aspects of the ecology and natural history of *paraergasilus rylovi* (copepoda, ergasilidae) parasitic in unionids of Finland. *Journal of parasitology* 90(5): 948–952.
- Smith, C., M. Reichard, P. Jurajda, and M. Przybylski. 2004. The reproductive ecology of the European bitterling (*Rhodeus sericeus*). *Journal of Zoology* 262: 107–124.
- Taskinen, J. 1998. Influence of trematode parasitism on the growth of a bivalve host in the field. *International Journal for Parasitology* 28: 599–602.
- Taskinen, J., and Saarinen, M. 1999. Increased Parasite Abundance Associated with Reproductive Maturity of the Clam *Anodonta piscinalis*. *The Journal of Parasitology* 85(3): 588.
- Taskinen, J., and M. Saarinen. 1999. Increased parasite abundance associated with reproductive maturity of the clam *Anodonta piscinalis*. *Journal of Parasitology* 85: 588–591.

Bibliography

- Taskinen, J., E. T. Valtonen, and T. Makela. 1994. Quantity of sporocysts and seasonality of two Rhipidocotyle species (Digenea: Bucephalidae) in *Anodonta piscinalis* (Mollusca: Bivalvia). *Int. J. Parasitol.*, 24: 877-886
- Toews, S., M. Beverly-Burton, and T. Lawrimore. 1993. Helminth and protist parasites of zebra mussels, *Dreissena polymorpha* (Pallas, 1771), in the Great Lakes region of southwestern Ontario, with comments on associated bacteria. *Can. J. Zool.*, 71: 1763-1766
- Zdun, V. I. 1961. Trematode larvae of freshwater molluscs from the Ukraine, Kiev: Vydvo (in Ukrainian)
- Zdun, V. I., V. K. Kiselene, A. Y. Karatayev, and G. E. Makarova. 1994. Parasites. In: *Freshwater Zebra Mussel Dreissena polymorpha* (Pall.) (Bivalvia, Dreissenidae). Systematics, Ecology, Practical Meaning, pp. 196-205. (J. I. Starobogatov, Ed.). Moscow: Nauka (in Russian)
- Zieritz, A., and Aldridge, D.C. 2011. Sexual, habitat-constrained and parasite-induced dimorphism in the shell of a freshwater mussel (*Anodonta anatina*, Unionidae). *Journal of Morphology* 272(11): 1365–1375.

Fresh Water Bivalve Bacteriology:

- Antunes, F., Hinzmann, M., Lopes-Lima, M., Machado, J., and Martins da Costa, P. 2010. Association Between Environmental Microbiota and Indigenous Bacteria Found in Hemolymph, Extrapallial Fluid and Mucus of *Anodonta cygnea* (Linnaeus, 1758). *Microbial Ecology* 60(2): 304–309.
- Chittick*, B., Stoskopf, M., Law, M., Overstreet, R., and Levine, J. 2001. Evaluation of potential health risks to Eastern Elliptio (*Elliptio complanata*) (Mollusca: Bivalvia: Unionida: Unionidae) and implications for sympatric endangered freshwater mussel species. *Journal of Aquatic Ecosystem Stress and Recovery* 9(1): 35–42.
- Fokin, S. I., Giamberini, L., Molloy, D. P., and bij de Vaate, A. 2003. Bacterial endocytobionts within endosymbiotic ciliates in *Dreissena polymorpha* (Lamellibranchia: Mollusca). *Acta Protozool.* 42:31-39.
- Fuller, S. L. H. 1974. Clams and mussels (Mollusca: Bivalvia). Pages 215-273 in *Pollution Ecology of Freshwater Invertebrates* (Hart, Jr. C. W. and Fuller, S. L. H., eds.). Academic Press, New York.
- Gu, J.-D., and Mitchell, R. 2002. Indigenous microflora and opportunistic pathogens of the freshwater zebra mussel, *Dreissena polymorpha*. *Hydrobiologia* 474(1): 81–90.
- Molloy, D. P., Mayer, D. A., Gaylo, M. J., Burlakova, L. E., Karatayev, A. Y., Presti, K. T., Sawyko, P. M., Morse, J. T., Paul, E. A. 2013. Non-target trials with *Pseudomonas fluorescens* strain CL145A, a lethal control agent of dreissenid mussels (Bivalvia: Dreissenidae). *Manag. Biol. Invasions* 4(1):71-79.
- Molloy, D. P., Mayer, D. A., Giamberini, L., and Gaylo, M. J. 2013. Mode of action of *Pseudomonas fluorescens* strain CL145A, a lethal control agent of dreissenid mussels (Bivalvia: Dreissenidae). *J. Invertebr. Pathol.* 113(1):115-121.
- Starliper, C.E. 2008. Recovery of a fish pathogenic bacterium, *Aeromonas salmonicida*, from ebonyshell mussels *Fusconaia ebena* using nondestructive sample collection procedures. *Journal of Shellfish Research* 27(4): 775–782.
- Starliper, C.E. 2009. Pathogens and diseases of freshwater mussels in the United States: studies on bacterial transmission and depuration. In *Bridging America and Russia with Shared Perspectives on Aquatic Animal Health. Proceedings of the Third Bilateral Conference between Russia and the United States.* pp. 12–20.
- Starliper, C.E., and Morrison, P. 2000. Bacterial pathogens contagion studies among freshwater bivalves and salmonid fishes. *Journal of Shellfish Research* 19(1): 251–258.
- Starliper, C.E., Neves, R.J., Hanlon, S., and Whittington, P. 2008. A survey of the indigenous microbiota (Bacteria) in three species of mussels from the Clinch and Holston rivers, Virginia. *Journal of Shellfish Research* 27(5): 1311–1317.
- Starliper, C.E., Powell, J., Garner, J.T., and Schill, W.B. 2011. Predominant Bacteria Isolated from Moribund *Fusconaia ebena* Ebonyshells Experiencing Die-Offs in Pickwick Reservoir, Tennessee River, Alabama. *Journal of Shellfish Research* 30(2): 359–366.
- Winters, A.D., Marsh, T.L., and Faisal, M. 2010. Bacterial Assemblages Associated with Zebra Mussel (*Dreissena polymorpha*) Populations in the Laurentian Great Lakes Basin (USA). *Journal of Shellfish Research* 29(4): 985–987.

Bibliography

Freshwater Bivalve Methodology

Histopathology:

- Bowmer, C. T. and M. van der Meer. 1991. Reproduction and Histopathological Condition in First Year Zebra Mussels (*Dreissena polymorpha*) From the Haringvliet, Volkerakmeer and Hollands Diep Basins. TNO Institute of Environmental Sciences Report R91/132, Delft, The Netherlands.
- Grizzle, J. M. and Brunner, C. J. 2007. Assessment of Current Information Available for Detection, Sampling, Necropsy, and Diagnosis of Diseased Mussels. Alabama Department of Conservation and Natural Resources Wildlife and Freshwater Fisheries Division, Montgomery, Alabama. 82 pp.
- McElwain, A., and Bullard, S.A. 2014. Histological atlas of freshwater mussels (Bivalvia, Unionidae): *Villosa nebulosa* (Ambleminae: Lampsilini), *Fusconaia cerina* (Ambleminae: Pleurobemini) and *Strophitus connasaugaensis* (Unioninae: Anodontini). *Malacologia* 57(1): 99–239.
- Morton, B. 1974. Studies on the biology of *Dreissena polymorpha*. VI. The occurrence of chronic pallial and tenidial inflammatory granulomas - the response to injury. *J. Invertebr. Pathol.*, 23: 106-113

Assessment:

- Burkhard, M.J., Leavell, S., Weiss, R.B., Kuehn, K., Valentine, H., Thomas Watters, G., and Wolfe, B.A. 2009. Analysis and cytologic characterization of hemocytes from freshwater mussels (*Quadrula* sp.). *Veterinary Clinical Pathology* 38(4): 426–436.
- Fritts, A.K., Peterson, J.T., Hazelton, P.D., Bringolf, R.B., and MacLatchey, D. 2015. Evaluation of methods for assessing physiological biomarkers of stress in freshwater mussels¹. *Canadian Journal of Fisheries and Aquatic Sciences* 72(10): 1450–1459.
- Gustafson, L.L., Stoskopf, M.K., Bogan, A.E., Showers, W., Kwak, T.J., Hanlon, S., and Levine, J.F. 2005. Evaluation of a nonlethal technique for hemolymph collection in *Elliptio complanata*, a freshwater bivalve (Mollusca: Unionidae). *Diseases of aquatic organisms* 65(2): 159–165.
- Newton T.J. and W.G. Cope. 2007. Biomarker responses of unionid mussels to environmental contaminants. Pages 257 to 284 in *Freshwater Bivalve Ecotoxicology*, J.L. Farris and J.H. Van Hassel, eds., SETAC Press, Pensacola, FL and Taylor & Francis, Boca Raton, FL.

Introduction to Omics:

- Eklblom R, Galindo J (2011) Applications of next generation sequencing in molecular ecology of non-model organisms. *Heredity* 107:1-15.
- Sardans J, Peñuelas J, Rivas-Ubach A (2011) Ecological metabolomics: overview of current developments and future challenges. *Chemoecology* 21:191-225.
- Suárez-Ulloa V, Fernández-Tajes J, Manfrin C, Gerdol M, Venier P, Eirín-López JM (2013) Bivalve omics: state of the art and potential applications for the biomonitoring of harmful marine compounds. *Mar Drugs* 11:4370-4389.

Unionid transcriptomics:

- Cornman RS, Robertson LS, Galbraith H, Blakeslee C (2014) Transcriptomic analysis of the mussel *Elliptio complanata* identifies candidate stress-response genes and an abundance of novel or noncoding transcripts. *PloS One* 9, e112420.
- Luo Y, Li C, Landis AG, Wang G, Stoeckel J, Peatman E (2014) Transcriptomic profiling of differential responses to drought in two freshwater mussel species, the giant floater *Pyganodon grandis* and the pondhorn *Unio merus tetralasmus*. *PloS One* 9, e89481.
- Robertson LS, Galbraith HS, Iwanowicz D, Blakeslee CJ, Cornman RS (2017) RNA sequencing analysis of transcriptional change in the freshwater mussel *Elliptio complanata* after environmentally relevant sodium chloride exposure. *Environ Toxicol Chem* 36:2352-2366.
- Wang R, Li C, Stoeckel J, Moyer G, Liu Z, Peatman E (2012) Rapid development of molecular resources for a freshwater mussel, *Villosa lienosa* (Bivalvia: Unionidae), using an RNA-seq-based approach. *Freshw Sci* 31:695-708.

Bibliography

Unionid metabolomics:

- Leonard JA, Cope WG, Barnhart MC, Bringolf RB (2014) Metabolomic, behavioral, and reproductive effects of the synthetic estrogen 17 α -ethinylestradiol on the unionid mussel *Lampsilis fasciola*. *Aquat Toxicol* 150:103-116.
- Leonard JA, Cope WG, Barnhart MC, Bringolf RB (2014) Metabolomic, behavioral, and reproductive effects of the aromatase inhibitor fadrozole hydrochloride on the unionid mussel *Lampsilis fasciola*. *Gen Comp Endocr* 206:213-226.
- Roznere I, Watters GT, Wolfe BA, Daly M (2014) Nontargeted metabolomics reveals biochemical pathways altered in response to captivity and food limitation in the freshwater mussel *Amblema plicata*. *Comp Biochem Physiol D* 12:53-60.
- Roznere I, Watters GT, Wolfe BA, Daly M (2017) Effects of relocation on metabolic profiles of freshwater mussels: metabolomics as a tool for improving conservation techniques. *Aquat Conserv* 27:919-926.

Conservation Implications of Omics:

- Allendorf FW, Hohenlohe PA, Luikart G (2010) Genomics and the future of conservation genetics. *Nat Rev Genet* 11:697-709.
- Corlett RT (2017) A bigger toolbox: biotechnology in biodiversity conservation. *Trends Biotechnol* 35:55-65.
- He X, Wilson CC, Wellband KW, Houde ALS, Neff BD, Heath DD (2015) Transcriptional profiling of two Atlantic salmon strains: implications for reintroduction into Lake Ontario. *Conserv Genet* 16:277-287.

Risk Assessment:

- Jakob-Hoff, R. M., S. C. MacDiarmid, C. Lees, P. S. Miller, D. A. Travis, and R. Kock. 2014. *Manual of Procedures for Wildlife Disease Risk Analysis*. Paris: OIE.
- NRC. 2003. "Risk Assessment: Evaluating Risks to Human Health and Safety." In *Occupational Health and Safety in the Care and Use of Nonhuman Primates*, Chapter 5. Washington: National Academies Press. <http://www.ncbi.nlm.nih.gov/books/NBK43454/>
- OIE, and IUCN. 2014. *Guidelines for Wildlife Disease Risk Analysis*, edited by R. Kock, W. B. Karesh, L. Skerratt, M. Hartley, and D. A. Travis. Paris: OIE.
- Miller, P. 2007. "Tools and Techniques for Disease Risk Assessment in Threatened Wildlife Conservation Programmes." *International Zoo Yearbook*, 41: 38–51.

Die-off:

- Ahlstedt, SA. 1989. Update of the Watts Bar Nuclear Plant preoperational monitoring of the mussel fauna in Upper Chickamauga Reservoir. Technical Report, Tennessee Valley Authority, Water Resources, Aquatic Biology Department.
- Blodgett, KD, Sparks RE. 1987. Analysis of a mussel die-off in pools 14 and 14 of the Upper Mississippi River. Aquatic biology technical report submitted to non-game check-off program, Illinois Department of Conservation.
- Dixon W. 2005. Disappearance of clams a sign of dying pond? Warwick Beacon, 4 Aug. 2005, www.buckeyebrook.org/news/warbeacon_clams.html.
- Fleming WJ, Augspurger TP, Alderman JA. 1995. Freshwater mussel die-off attributed to anticholinesterase poison. *Environmental Toxicology and Chemistry*. 14, 877–879. 10.1002/etc.5620140520
- Fralely SJ, Simmons JW. 2006. An assessment of selected rare mussel populations in Western North Carolina following extraordinary floods of September 2004. Technical report, North Carolina Wildlife Resources Commission.
- Gomez I, Araujo R. 2008. Channels and ditches as the last shelter for freshwater mussels: the case of *Margaritifera auricularia* and other naiads inhabiting the mid Ebro River Basin, Spain. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18: 658-670.
- Holcomb J. 2017. Lake Victor Mussel Kill Report. Technical report, Florida Fish and Wildlife Conservation Commission.
- Hubbs, D. 2001. 2001 Tennessee statewide commercial mussel report. Technical report, Tennessee Wildlife Resources Agency.

Bibliography

- Hubbs, D. 2007. 2007 Tennessee statewide commercial mussel report. Technical report, Tennessee Wildlife Resources Agency.
- Jones JW, Neves RJ. 2005. Final Report: A status survey of freshwater mussel populations in the Upper North Fork Holston River, Virginia. Technical report, Virginia Cooperative Fish and Wildlife Research Unit.
- Kat, P. W. 1982. Shell dissolution as a significant cause of mortality for *Corbicula fluminea* (Bivalvia: Corbiculidae) inhabiting acidic waters. *Malacological Review* 15: 129–134.
- Levine JF, Salger S, Borst L, Law M, Eads C, Dykstra M, Osburn C, Gangloff M, Sumner S, Saul B. 2015. Investigation of factors contributing to the decline in the nutritional health of *Alasmidonta raveneliana* in the Little Tennessee River, Franklin, North Carolina. Technical report submitted to NCDOT, project 2013-2013-37.
- Miller JR, Mackin G. 2013. Concentrations, sources, and potential ecological impacts of selected trace metals on aquatic biota within the Little Tennessee River Basin, North Carolina. *Water, Air, & Soil Pollution* 224:1613.
- Miller JR. 2016. Potential ecological impacts of trace metals on aquatic biota within the Upper Little Tennessee River Basin, North Carolina. *AIMS Environmental Science* 3(3): 305-325.
- Sasson A. 2017. Summary of 2016 Big Darby Creek mussel die-off. Technical Report, The Nature Conservancy.
- Southwick RI, Loftus AJ. 2003. Investigation and Monetary Values of Fish and Freshwater Mussel Kills, Special Publication 30. Bethesda, MD: American Fisheries Society.
- Southwick, R.I., and A.J. Loftus, editors. 2017. Investigation and monetary values of fish and freshwater mollusk kills. American Fisheries Society, Special Publication 35, Bethesda, Maryland.
- Sparks RE, Blodgett KD, Durham L, Horner R. 1990. Determination Whether the Causal Agent for Mussel Die-offs in the Mississippi River is of Chemical or Biological Origin. Final Report, ILENR/RE-WR 90/09, Illinois Department of Energy and Natural Resources, Springfield, IL.

Attendee List

Alfjorden	Anders	National Veterinary Institute (SVA)	anders.alfjorden@sva.se
Aloisi	Doug	Genoa National Fish Hatchery- USFWS	doug_aloisi@fws.gov
Baran Dagendesh	Angela	Genoa National Fish Hatchery- USFWS	angela_baran_dagendesh@fws.gov
Boedeker	Nancy	Indiana Department of Natural Resources	nboedeker@dnr.in.gov
Bradley	Megan	USFWS- Genoa National Fish Hatchery	megan_bradley@fws.gov
Buntin	Michael	Alabama Aquatic Biodiversity Center	michael.buntin@dcnr.alabama.gov
Butitta	Vince	Center for Limnology	vincent.butitta@wisc.edu
Byrne	Caitlin	OSU	byrne.88@osu.edu
Carey	Caitlin	Virginia Tech, CMI/FiW	cscarey@vt.edu
Carlson	Brian	AllStar Ecology	brian@allstarecology.com
Carnegie	Ryan	Virginia Institute of Marine Science	carnegie@vims.edu
Carney	Joseph	Lakehead University, Biology Department	jcarney@lakeheadu.ca
Castrillo Arias	Pedro Antonio	Universidade de Santiago de Compostela	pedroantonio.castrillo@usc.es
Chance	Stephanie	USFWS - TN Ecological Services Field Office	stephanie_chance@fws.gov
Christian	Alan	University of Massachusetts Boston	alan.christian@umb.edu
Ciparis	Serena	US Fish and Wildlife Service	serena_ciparis@fws.gov
Clayton	Janet	WV Division of Natural Resources	janet.l.clayton@wv.gov
Colletti	Sarah	VA Dept. of Game and Inland Fisheries	sarah.colletti@dgif.virginia.gov
Cope	W. Gregory	North Carolina State University	greg_cope@ncsu.edu
Dahlberg	Angelique	St. Croix River Association	angeliqued@scramail.com
Dare	Jason	Dare Ecosystem Management	jason_dare@hotmail.com
Davis	Ben	USFWS	benjamin_davis@fws.gov
DeMartini	Jessi	Forest Preserve Dist. DuPage County Il.	jdemartini@dupageforest.org
Dorman	Rebecca	USGS, Columbia Environmental Research Center	rdorman@usgs.gov
Douglas	Barbara	USFWS	barbara_douglas@fws.gov
Douglass	Sarah	Illinois Natural History Survey	sabales@illinois.edu
Dunn	Christopher	UW-Madison	cddunn2@wisc.edu
Dunn	Heidi	EcoAnalysts, Inc.	hdunn@ecologicalspecialists.com
Eckert	Nathan	US Fish & Wildlife Service	nathan_eckert@fws.gov
Erickson	Sara	U.S. Fish & Wildlife Service	sara_erickson@fws.gov
Everhart	Michael	West Virginia Division of Natural Resources	mike.e.everhart@wv.gov
Faiman	Scott	MDC	scott.faiman@mdc.mo.gov
Finney	Sam	USFWS	sam_finney@fws.gov
Ford	David	Ecological Specialists, Inc.	dford@ecologicalspecialists.com
Fritts	Andrea	U.S. Geological Survey	afritts@usgs.gov
Ganser	Alissa	Virginia Tech	alissamganser@gmail.com
Glidewell	Elizabeth	(CTUIR) Confederated Tribes of the Umatilla Indian Reservation	ElizabethGlidewell@CTUIR.org
Goldberg	Tony	University of Wisconsin-Madison	tony.goldberg@wisc.edu
Grossman	Emily	EcoAnalysts, Inc.	egrossman@ecologicalspecialists.com
Haag	Wendell R	U.S. Forest Service	whaag@fs.fed.us
Haas	Jeremiah	Exelon	jeremiah.haas@exeloncorp.com
Halmbacher	Jacquelyn	The Ohio State University	halmbacher.2@osu.edu
Hampton	Dave	FWS	dave_hampton@fws.gov
Hazelton	Peter	Massachusetts Division of Fisheries & Wildlife	peter.hazelton@state.ma.us
Hern	Tyler	United States Fish and Wildlife Service	tyler_hern@fws.gov

Attendee List

Hoch	Rachael	North Carolina Wildlife Resources Commission	rachael.hoch@ncwildlife.org
Intihar	Jim	Forest Preserve District of DuPage County	jintihar@dupageforest.org
Jackson	Katelyn	EA Engineering, Science, and Technology, Inc, PBC	kjackson@eaest.com
Jenkinson	John	FMCS Member	jjjenkinson@hotmail.com
Johnson	Jennifer	Michigan DNR Fisheries	johnsonj17@michigan.gov
Kenney	Aleshia	USFWS	aleshia_kenney@fws.gov
Kern	Morgan	Missouri State University	mkern03@gmail.com
Kitchel	Lisie	WDNR - BER	lisie.kitchel@wisconsin.gov
Knowles	Susan	National Wildlife Health Center	sknowles@usgs.gov
Kunz	James	Columbia Environmental Research Center	jkunz@usgs.gov
Lane	Tim	Virginia Department of Game and Inland Fisheries	tim.lane@dgif.virginia.gov
Lavictoire	Louise	Missouri State University	louiselavictoire@hotmail.co.uk
Lawlis	Clarissa	Lewis Environmental Consulting, LLC	clarissa_lawlis@yahoo.com
Lennon	Tiernan	USFWS	tiernan_lennon@fws.gov
Leis	Eric	USFWS Midwest Fisheries Center	eric_leis@fws.gov
Lewis	Chad	Lewis Environmental Consulting, LLC	lewis_environmental@yahoo.com
Lewis	Teresa	USFWS Midwest Fisheries Center	teresa_lewis@fws.gov
Limpers	Joe	Forest Preserve District of DuPage County	jlimpers@dupageforest.org
Matthews	Mickey	Arkansas Dept of Transportation	mickey.matthews@ahd.ar.gov
Maynard	Amy	Virginia Department of Game and Inland Fisheries	amy.maynard@dgif.virginia.gov
Mazzacano	Celeste	CASM Environmental, LLC	cmazzacano@gmail.com
McComb	Sophia	Sycamore High School Watershed	mccombsoph@gmail.com
McCormick	Madeline	Sycamore High School/Student	madeline6520@gmail.com
McElwain	Andrew	SUNY Oswego	andrew.mcelwain@oswego.edu
McMurray	Stephen	MO Dept. of Conservation	stephen.mcmurray@mdc.mo.gov
Miller	Phil	Department of Veterinary Population Medicine	
Millsap	Deborah	US FWS	deborah_millsap@fws.gov
Minerich	Ben	Minnesota Zoo	ben.minerich@state.mn.us
Molloy	Daniel	Molloy & Associates, LLC	dan@danielpmolloy.com
Monroe	Emy	US Fish & Wildlife Service	emy_monroe@fws.gov
Morrison	Patricia	USFWS Retired	pearlymussel@gmail.com
Mosier II	Dan	Kansas Dep. of Wildlife, Parks & Tourism	dan.mosier@ks.gov
Moss	Lindsey	TRC	lmoss@trcsolutions.com
Nemeth	Doug	Idaho Fish and Wildlife Conservation Office	douglas_nemeth@fws.gov
Newton	Teresa	USGS	tnewton@usgs.gov
Ohlman	Lindsay	University of Nebraska	lohlman@unl.edu
Ostby	Brett	Daguna Consulting, LLC	ptychobranchus@gmail.com
Parsons	Edward	Greensboro Science Center	eparsons@greensboroscience.org
Patterson	Matthew	USFWS	matthew_patterson@fws.gov
Penn	Michael	USFWS - Lamar Fish Health Center	michael_penn@fws.gov
Phelps	Nick	Minnesota Aquatic Invasive Species Research Center	phelp083@umn.edu
Primus	Alex	Department of Veterinary Population Medicine	primu012@umn.edu
Riccardi	Nicoletta	CNR Institute of Ecosystem Study	n.riccardi@ise.cnr.it
Richard	Jordan	US Fish and Wildlife Service	jordan_richard@fws.gov

Attendee List

Attendee List

Rosenberger	Amanda	USGS Missouri Cooperative Fish and Wildlife Research Unit	rosenbergera@missouri.edu
Roznere	Ieva	The Ohio State University	roznere.1@osu.edu
Runstrom	Ann	U.S. Fish and Wildlife Service	ann_runstrom@fws.gov
Sasson	Anthony	The Nature Conservancy	asasson@aol.com
Schmuecker	Sara	U.S. Fish and Wildlife Service	sara_schmuecker@fws.gov
Scoggin	Dan	Ecoanalysts, Inc.	scogdw@hotmail.com
Seagraves	Sara	US Fish and Wildlife Service	sara_seagraves@fws.gov
Secrist	Zebulin	Dept. of Natural Resources	zeb.secrist@state.mn.us
Sietman	Bernard	Minnesota DNR	bernard.sietman@state.mn.us
Sneed	Lesley	Kentucky State University	lesley.sneed@kysu.edu
Taskinen	Jouni	University of Jyväskylä	jouni.k.taskinen@jyu.fi
Tiemann	Jeremy	Illinois Natural History Survey	jtiemann@illinois.edu
Urbańska	Maria	Poznań University of Life Sciences, Institute of Zoology	urbanska@up.poznan.pl
Vaughn	Chris	San Antonio River Authority	cvvaughn@sara-tx.org
Waller	Diane	USGS-UMESC	dwaller@usgs.gov
Watson	Brian	VA Dept of Game & Inland Fisheries	brian.watson@dgif.virginia.gov
Weinzinger	Jesse	Wisconsin DNR	jesse.j.weinzinger@gmail.com
Wengstrom	Niklas	University of Gothenburg	niklas.wengstrom@bioenv.gu.se
Wolf	Tiffany	Department of Veterinary Population Medicine	wolfx305@umn.edu
Woolnough	Daelyn	Central Michigan University	daelynw@gmail.com

From the City of La Crosse and the 2018 Workshop Planning Committee:

***Thank you for making the
2018 Workshop a Success!***

